

SOQUEL DEMONSTRATION
STATE FOREST

DRAFT

GENERAL FOREST
MANAGEMENT PLAN

**CALIFORNIA DEPARTMENT OF FORESTRY
AND FIRE PROTECTION**

SOQUEL, CALIFORNIA

August, 2011

Richard Sampson, Acting Forest Manager
Edgar Orre, Assistant Forest Manager

SDSF ADVISORY COMMITTEE

California Department of Fish and Game
Stewards of Soquel Forest
County of Santa Cruz Planning Department
Neighborhood Representative
California Board of Forestry and Fire Protection
The Nature Conservancy
Registered Professional Forester
California Department of Parks and Recreation
Resource Conservation District of Santa Cruz County

KEN PIMLOTT
ACTING DIRECTOR
CALIFORNIA DEPARTMENT OF FORESTRY
AND FIRE PROTECTION

EDMUND G. BROWN, JR.
GOVERNOR
STATE OF CALIFORNIA

JOHN LAIRD
SECRETARY FOR RESOURCES
THE NATURAL RESOURCES AGENCY

TABLE OF CONTENTS

INTRODUCTION	1
CALIFORNIA'S DEMONSTRATION STATE FOREST SYSTEM.....	1
THE GENERAL FOREST MANAGEMENT PLAN	1
FUTURE PLANNING.....	2
MANAGEMENT GOALS	3
DEMONSTRATION AND EDUCATION	3
TIMBER MANAGEMENT.....	4
RESEARCH	4
RESOURCE PROTECTION AND ENHANCEMENT.....	4
RECREATION	5
PUBLIC COMMENTS AND CONCERNS.....	5
PROPERTY DESCRIPTION	6
LOCATION AND BOUNDARIES	6
ADJACENT OWNERSHIP.....	6
HISTORY OF OWNERSHIP.....	6
CLIMATE.....	7
SOILS AND GEOLOGY.....	7
WATER RESOURCES	10
ZONING AND GENERAL PLAN	10
CHARACTERISTIC FEATURES	10
ADMINISTRATION	12
THE LEASE	12
CDF ADMINISTRATION	12
FUNDING AND TAXES	13
ADVISORY COMMITTEE	14
SAFETY	15
LAW ENFORCEMENT.....	15
ACCESS NEEDS.....	16
LAND ACQUISITION PRIORITIES	16
COOPERATION WITH ADJACENT LANDOWNERS	16
COOPERATION WITH OTHER AGENCIES AND INSTITUTIONS.....	16
PUBLIC COMMENTS AND CONCERNS.....	16
BIOTA	19
INTRODUCTION	19
HABITAT COMMUNITIES.....	19
FUNGAL RESOURCES	21
PLANT SPECIES OF CONCERN	22
SPECIAL-STATUS WILDLIFE SPECIES	22
PUBLIC COMMENTS AND CONCERNS.....	24
MANAGEMENT GUIDELINES.....	25
PLANNED ACTIONS.....	25

FISHERIES.....	27
INTRODUCTION	27
HABITAT AND RESOURCE CONSIDERATIONS.....	29
AQUATIC INVERTEBRATES.....	30
PUBLIC COMMENTS AND CONCERNS.....	31
MANAGEMENT GUIDELINES.....	32
PLANNED ACTIONS.....	32
DEMONSTRATION AND EDUCATION	34
INTRODUCTION	34
DEMONSTRATION PROGRAM	34
FORESTRY EDUCATION.....	36
INTERPRETATION.....	37
VOLUNTEER PROGRAM.....	38
TIMBER MANAGEMENT.....	39
INTRODUCTION	39
HARVEST HISTORY	39
CONIFER VOLUME INVENTORY	40
GROWTH	41
SILVICULTURE.....	42
OLD-GROWTH AND LATE-SUCCESSION MANAGEMENT.....	45
HARVEST SYSTEMS	47
HARVEST PLAN.....	47
TIMBER OPERATIONS MONITORING.....	53
TIMBER MARKETS	53
HARDWOOD MANAGEMENT.....	54
DEMONSTRATION AND RESEARCH PROGRAMS	55
MINOR FOREST PRODUCTS	56
PUBLIC COMMENTS AND CONCERNS.....	56
MANAGEMENT GUIDELINES.....	57
PLANNED ACTIONS.....	58
RESEARCH.....	60
INTRODUCTION	60
COMPLETED PROJECTS	60
RESEARCH PRIORITIES	62
PUBLIC COMMENTS AND CONCERNS.....	63
MANAGEMENT GUIDELINES.....	63
PLANNED ACTIONS.....	63
RECREATION	65
INTRODUCTION	65
THE RECREATION STUDY	65
EXISTING FACILITIES.....	68
PUBLIC COMMENTS AND CONCERNS.....	70
MANAGEMENT GUIDELINES.....	70

PLANNED ACTIONS.....	71
RESOURCE PROTECTION.....	73
FIRE PROTECTION.....	73
PUBLIC COMMENTS AND CONCERNS.....	77
PLANNED ACTIONS.....	78
FOREST PEST MANAGEMENT.....	78
PLANNED ACTIONS.....	85
WATERSHED ASSESSMENT.....	87
PHYSIOGRAPHY OF THE WATERSHED.....	87
BENEFICIAL USES.....	87
LAND USE HISTORY.....	88
CUMULATIVE WATERSHED EFFECTS ASSESSMENT.....	89
MONITORING AND ENHANCEMENT.....	93
PUBLIC COMMENTS AND CONCERNS.....	93
MANAGEMENT GUIDELINES.....	93
PLANNED ACTIONS.....	94
ARCHAEOLOGY.....	95
INTRODUCTION.....	95
PREHISTORIC SITES.....	95
HISTORIC SITES.....	96
PUBLIC COMMENTS AND CONCERNS.....	97
MANAGEMENT GUIDELINES.....	97
PLANNED ACTIONS.....	97
ROADS AND OTHER IMPROVEMENTS.....	99
SDSF FACILITIES.....	99
PUBLIC ACCESS ROADS.....	100
ROAD BUILDING AND MAINTENANCE.....	100
PUBLIC COMMENTS AND CONCERNS.....	101
MANAGEMENT GUIDELINES.....	102
PLANNED ACTIONS.....	102
REFERENCES.....	104
LITERATURE.....	104
PERSONAL COMMUNICATION.....	110
APPENDICES.....	111
APPENDIX A: TEXT OF ASSEMBLY BILL 1965.....	111
APPENDIX B: SPECIES LISTS FOR SDSF.....	113
FAUNA OF SDSF.....	113
FLORA OF SDSF.....	118
FUNGI OF SDSF.....	125
APPENDIX C: MONITORING PLAN.....	127

LIST OF FIGURES

Figure 1.	Locator map for SDSF.....	133
Figure 2.	SDSF and adjacent ownerships.....	134
Figure 3.	Locations of soil series in SDSF.....	135
Figure 4.	Vegetation Type Map for SDSF.....	136
Figure 5.	Site Class Map for SDSF.....	137
Figure 6.	Diameter class distribution for SDSF.....	43
Figure 7.	Old-growth and late-succession management areas.....	138
Figure 8.	Preliminary logging plan.....	139
Figure 9.	Decadel harvest, decadel growth, and inventory of available acres by planning periods.....	51
Figure 10.	Projected annual conifer growth and harvest on SDSF timberlands with full or limited availability.....	51
Figure 11.	Projected aggregate conifer stocking on SDSF timberlands with full or limited availability.....	52
Figure 12.	Timber market stumpage prices.....	54
Figure 13.	Estimated basal area for SDSF's hardwood tree species.....	55
Figure 14.	Recreational trails of SDSF and potential campground location.....	140
Figure 15.	Proposed fire defense improvements.....	141
Figure 16.	Annual Peak Streamflow for Soquel Creek.....	87
Figure 17.	Drivable roads of SDSF and surrounding area.....	142

LIST OF TABLES

Table 1.	Soil types and associated erosion hazard of SDSF.....	9
Table 2.	Special-status wildlife species with potential to occur at Soquel Demonstration State Forest.....	23
Table 3.	SDSF 2006 Forest Resources Inventory Summary	41
Table 4.	Individual redwood and Douglas-fir tree characteristics.....	47
Table 5.	Soquel DSF forestland availability.....	50
Table 6.	Decadel harvest, growth and inventory.....	50
Table 7.	Forest pests of the Soquel Creek watershed.....	79
Table 8.	Drivable road segments of SDSF.....	101

INTRODUCTION

CALIFORNIA'S DEMONSTRATION STATE FOREST SYSTEM

The demonstration state forest system of the California Department of Forestry and Fire Protection (CAL FIRE) was established in the mid-1940's to furnish local needs of research, demonstration, and education related to forest management. Currently, the demonstration state forest (DSF) system encompasses over 71,000 acres of land in the form of eight state forests. DSFs are healthy, living forests which demonstrate conservation and protection of wildlife, fisheries, vegetation, soil, and watershed resources as well as sustained-yield forest management activities.

The Soquel Demonstration State Forest (SDSF), when established in 1990, was the first addition to the DSF system in over 40 years. SDSF contains approximately 2,681 acres of coast redwood and mixed evergreen forest types. Former Assemblyman Sam Farr authored SDSF's enabling legislation, Assembly Bill 1965 of 1987 (now Public Resources Code (PRC) Sections 4660-4664), which provided for the protection and preservation of the SDSF as an intensively managed educational and research forest, and contained special provisions for the use of SDSF, including a limited amount of commercial timber operations on the property within the SDSF in order to provide funds for the maintenance and operation of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Section 4660 on SDSF. AB 1965 is reprinted in Appendix A of this plan.

SDSF was formally transferred from its interim managers, The Nature Conservancy (TNC), to CAL FIRE on July 13, 1990. Former CAL FIRE Director Harold Walt, former Assemblyman Sam Farr, a representative of State Controller Gray Davis, and Steve Johnson of TNC dedicated the Forest, emphasizing the nature and purpose of this addition to the DSF system.

SDSF was the first of California's demonstration state forests to have an advisory committee formed to assist the Department in planning future management of the Forest. The Advisory Committee, also required by AB 1965, met monthly during the planning process to facilitate the creation of the original 1998 General Forest Management Plan. The Advisory Committee will be reconvened and will play a vital role in reviewing the revisions to this updated version of the plan.

THE GENERAL FOREST MANAGEMENT PLAN

Following the acquisition of the SDSF property (see the Administration chapter), TNC created an interim management plan for what they called the Soquel Creek Forest. Recognized as a temporary plan, TNC's document provided direction for current and future management decisions involving SDSF. The 1998 General Forest Management Plan incorporated elements of the TNC plan and information from other sources. It was developed with input from the public and resource professionals. Public workshops were held to obtain feelings, opinions, and factual information about the management of SDSF's forest resources. Furthermore, individuals representing many interests contributed data, publications, and personal knowledge for consideration through conversation with Forest staff. Public comments and concerns relating to

various subjects are summarized in each chapter. A supplemental document, titled Public Input to the Soquel Demonstration State Forest General Management Plan, presents all input received during the development of the draft plan which was completed in October, 1993.

Local resource professionals contributed a significant amount to the 1998 plan. Knowledgeable individuals served on the Advisory Committee, provided factual information about various resources, and composed elements of the plan itself. Without their assistance and experience, this management plan could not have been written.

The environmental impacts of the 1998 management plan were thoroughly investigated by Jones and Stokes Associates, Inc. in a program environmental impact report (EIR) which was completed in compliance with the California Environmental Quality Act (CEQA). A primary objective of this program EIR is to identify mitigation measures to reduce or avoid adverse environmental impacts that could result from implementation of any SDSF projects. As required by CEQA, mitigation measures identified from this EIR have been incorporated into a Monitoring Plan located in Appendix C.

FUTURE PLANNING

The Board of Forestry and Fire Protection (the Board) approved the original General Forest Management Plan on May 18, 1998. In 2003, the Board reviewed and reauthorized the plan. No revisions were made at that time.

As indicated throughout this plan, SDSF staff continue to formulate more specific management guidelines and planned actions. This revision of the plan includes new studies and the results of monitoring and research regarding the management of components such as , fisheries, wildlife, watershed, and timber. .

This SDSF General Forest Management Plan will be in effect until it is either amended or a new plan is adopted in accordance with the procedures prescribed in PRC Section 4663. Working with the Advisory Committee, CAL FIRE will reexamine the General Forest Management Plan every five years and determine whether any changes are necessary or desirable. This plan embodies the legislative intent of PRC Sections 4660-4664, and any subsequent amendments of this plan or any new plan must be consistent with the interpretations of PRC Sections 4660-4664 contained in this plan except to the extent, if any, that subsequent legislation changes that intent. If changes are desired, the changes will be developed by CAL FIRE and presented to the Advisory Committee for consideration at one or more public meetings. The changes shall be approved by the Advisory Committee prior to adoption by the Board. Any changes will be subject to environmental review as provided by the California Environmental Quality Act.

MANAGEMENT GOALS

SDSF's management goals represent a combination of legislation, policy, and public input. PRC Sections 4660-4664, the enabling legislation for Soquel Demonstration State Forest, is the preeminent authority with regard to the management of the SDSF. Consistent with the objectives of that legislation to protect and preserve SDSF as an intensively managed, multifaceted research forest and to the extent not in conflict with that enabling legislation, the SDSF will be managed in accordance with the state forest system legislation (PRC Sections 4631-4658) and Board of Forestry policy.

Public Resources Code Section 4660 states that the intent of the Legislature in establishing the Soquel Demonstration State Forest is to provide an environment that will do all of the following:

- * Provide watershed protection for local communities and base-line monitoring and studies of the hazards, risks, and benefits of forest operations and watersheds to urban areas.
- * Provide public education and examples illustrating compatible rural land uses, including sustained yield timber production, as well as the historic development of timbering and forestry machinery, within the context of local community protection and nearby pressures.
- * Provide a resource for the public, environmental groups, elected officials, environmental planners, the educational community, and the media as an open environment for the inspection and study of environmental education, forestry practices, and effects thereof.
- * Protect old-growth redwood trees.

As is common in legislation, these objectives contain many potential conflicts and will require trade offs in implementation.

Section 4661 further states that CAL FIRE may permit a limited amount of commercial timber harvesting in order to provide the funds needed for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF.

Below is a listing of SDSF's general management goals which elaborate on the legislative intent. Other subjects and greater detail relating to the topics listed here can be found throughout the following chapters of this General Forest Management Plan.

DEMONSTRATION AND EDUCATION

1. Conduct innovative demonstrations and education in forest management including silviculture, habitat diversity, logging methods, hydrology, resource protection, and recreation.

2. Provide forestry education opportunities for the public, forest landowners, the educational community, the media, natural resource professionals, and environmental groups.
3. Develop interpretive resources to help Forest visitors understand the various coast redwood forest communities and the basics of forest land management.
4. Establish a volunteer program to assist forest staff in providing forestry interpretation for visitors.
5. Design and construct a Forestry Education Center to serve as the Forest's focal point for demonstration and education activities.

TIMBER MANAGEMENT

1. Demonstrate sustained-yield with examples of timber harvesting at a level that is compatible with rural land use in Santa Cruz County and recreational use of SDSF and promotes forest health, watershed protection, wildlife, and fisheries values as well as aesthetic enjoyment.
2. Protect old-growth redwood and Douglas-fir trees and recruit additional late-successional forest stands.
3. Incorporate demonstration, research, and restoration objectives into timber management activities whenever possible.
4. Study hardwood stand management alternatives including modification to enhance wildlife habitat, utilization for various forest products, and conversion to softwood timber stands consistent with the legislative goals of PRC Sections 4660-4664.

RESEARCH

Conduct research in forestry and natural resource management, including the benefits and risks of forest operations in watersheds close to urban areas.

Serve as a laboratory for in-house projects and encourage research by other agencies, interest groups, and educational institutions.

Disseminate information obtained from the State Forest to appropriate individuals in an effective and timely manner.

RESOURCE PROTECTION AND ENHANCEMENT

1. Protect, restore, and enhance the significant natural values of the Soquel Demonstration State Forest.
2. Provide watershed protection and conduct baseline studies and monitoring of

hydrological resources.

3. Demonstrate fire protection using a coordinated fire prevention and control system which includes education and enforcement of fire prevention guidelines, Forest patrol, vegetation management, fuelbreak construction, pre-attack strategies, and suppression tactics.
4. Improve fisheries and wildlife habitat to foster healthy populations and promote biodiversity.
5. Monitor and study controls for various forest pests using Departmental and outside specialists.
6. Study all significant archaeological and historical features and protect them during all management activities.
7. Conserve soil resources by reducing erosion resulting from flooding, earthquakes, logging activities, roads, and trails.

RECREATION

1. Provide for recreational opportunities which are oriented toward foot, bicycle, and equestrian traffic and include trails, roads, and picnic areas. Limited camping may be permitted in the future.
2. Integrate recreation management, forestry education, resource protection and examples of timber harvesting so as to demonstrate how they can be compatible.
3. Control fishing, the use of motorized vehicles, shooting, and hunting to provide for public safety and forest protection.

PUBLIC COMMENTS AND CONCERNS

Throughout the planning process, members of the public have indicated that adherence to SDSF's enabling legislation, AB 1965, is legally required. Neighbors and visitors of the Forest have communicated that management goals and actions should abide by the written legislation. CAL FIRE intends to abide by this plan and to act consistently with the intention of the enabling legislation as expressed in this plan.

PROPERTY DESCRIPTION

LOCATION AND BOUNDARIES

Soquel Demonstration State Forest is located near the center of Santa Cruz County, California, approximately eight miles northeast of the city of Santa Cruz (Figure 1). Positioned in the southern portion of the Santa Cruz Mountains, SDSF is eighteen air miles south of San Jose and within a two-hour drive of the San Francisco and Oakland metropolitan areas. Access to the property is via State Highway 1 or 17 and local county roads. The entrance to the Forest is from Highland Way, a county road in the Santa Cruz Mountains that connects State Highway 17 with Watsonville. Virtually all of the Forest's 2,681 acres are located within the East Branch of Soquel Creek watershed.

SDSF's boundaries were originally established by metes and bounds rather than the more familiar township and range system. Formerly part of the Soquel Augmentation Rancho (a Mexican land grant), this area has always been defined differently than the land which surrounds it. Some of the corners listed in the survey, prepared by George Dunbar of Dunbar Land Surveys, have been verified and are in place. The Santa Cruz County parcel numbers for the Forest are 098-101-04; 098-161-06; 098-351-01; and 099-181-02, 03, 04, 06.

ADJACENT OWNERSHIP

SDSF is bordered by both state and private property (Figure 2). The Forest Of Nisene Marks State Park borders the State Forest for three and one-half miles along Santa Rosalia Ridge to the south. Approximately three-hundred-forty acres directly east of the Forest boundary are owned by Roger and Michelle Burch. This land is managed by Redwood Empire and includes the main entrance and parking area for the Forest off Highland Way. To the north and west, the adjacent ownerships are private rural-residential parcels, including the large holding of Spanish Ranch. Most of these parcels range in size from 1 to 80 acres. On the southwest border is the property containing the Olive Springs Quarry, owned by the CHY Company.

With the considerable amount of private property surrounding the Forest, public access is currently limited. The only undisputed public access points into the Forest are from Highland Way and The Forest of Nisene Marks State Park.

HISTORY OF OWNERSHIP

Prior to the arrival of Europeans, the Costanoan (also known as Ohlone) Indians inhabited the area. In the mid-1800's, the title to the 32,000-acre Soquel Augmentation Rancho was awarded to Martina Castro de Depeaux viuda de Lodge viuda de Cota, the daughter of a Spanish Colonial soldier. SDSF was contained within the rancho, and Martina gave this portion to her daughter, Antonia Lodge de Peck. Frederick A. Hihn, a German-born entrepreneur, was able to acquire portions of the Soquel Augmentation through a discrepancy in legal title. He was particularly interested in Lodge de Peck's parcel and purchased it in 1863.

In the 1880's, Hihn established the Valencia-Hihn Company and began selectively logging the old-growth redwood on his lands to produce shingles, posts, and rails. Upon his death in 1913, Hihn's heirs assumed management of his lands and continued to harvest the area. In 1924, the Valencia-Hihn Company sold their land to the Monterey Bay Redwood Company (MBRC). The MBRC owned the State Forest property for 37 years and performed extensive harvest in the 1920's and '30's. They sold their property to the Glenco Forest Products Company of Sacramento in 1961, which later changed its name to the CHY Company. Eighteen years later, in 1979, CHY sold the State Forest portion of their land to the Pelican Timber Company. Additional details about the history of the Forest can be found in Archaeological and Historical Survey of Soquel Demonstration State Forest (Dillon, 1992).

In 1988, Pelican was involved in a debt-for-nature land swap with the State of California and the Bank of America (see the Administration chapter for more details). A result of this land swap was the creation of SDSF as authorized by former Assemblyman Sam Farr's Assembly Bill 1965. The Nature Conservancy acted as the interim managers of the Forest until its transfer to CAL FIRE in 1990.

CLIMATE

The climate of the Santa Cruz Mountains is Mediterranean, characterized by dry, warm summers and wet, cool winters. SDSF is usually cool and damp because of the dense canopy of forest vegetation and its location on a north-facing slope. The average minimum January temperature is 38 degrees Fahrenheit, and the average maximum July temperature is 76 degrees Fahrenheit.

Most of the precipitation in the area occurs from November through April. The average annual rainfall for the East Branch of Soquel Creek is 44 inches (Linsley et al., 1992). At elevations above 2,000, snowfall occurs about every other year and averages less than five inches total.

During the late spring and early summer months, Santa Cruz County often has foggy or cloudy skies. In the Forest, this is generally limited to early morning and late evening hours. Winds generally blow from the west or southwest (onshore) and are mild to moderate throughout the year. Strong winds, however, come in with winter storms and are strongest at higher elevations. Pressure gradients inland may occasionally cause strong northeasterly winds to occur.

SOILS AND GEOLOGY

SOIL TYPES

The parent material of soils found in SDSF is primarily sedimentary and consists of fine and coarse-grained sandstone, consolidated shale, weathered mudstone, and siltstone. Schist and intrusions of granitic rock are also present. There are nine soil series which developed from these parent materials (Table 1 and Figure 3). They are all deep and well-drained soils except for the Maymen Stony Loam which is a shallow, well-drained soil. According to the Natural Resources Conservation Service (US Department of Agriculture, 1980), most of the soils support watershed, recreation, and wildlife resources. Five of the soils (Ben Lomond, Felton, Lompico, Nisene, and Aptos) also support timber production, with the primary species being coast

redwood and Douglas-fir.

GEOLOGIC ACTIVITY

In 1992, a detailed geologic study was completed by the California Geologic Survey (Manson and Sowma-Bawcom, 1992). This investigation resulted in a report which focuses on the process and degree of instability in both the State Forest and surrounding areas. The report, titled Geology, Slope Stability, and Earthquake Damage in Soquel Demonstration State Forest, includes maps of general geologic and geomorphic characteristics, landslide features (indicating the relative degree of stability), stream orders, roads to be considered for abandonment, and Alquist-Priolo Special Studies Zones. (Alquist-Priolo Special Studies Zones are areas along traces of the San Andreas Fault where geologic investigations are required prior to development.)

SDSF is seismically very active. The San Andreas Fault runs through the northeastern boundary and along the East Branch of Soquel Creek to the mouth of Ashbury Gulch, where it turns north. The Zayante Fault, part of the San Andreas Rift Zone, runs through the southwest edge of the Forest. The epicenter of the 1989 Loma Prieta Earthquake was located approximately 2 miles south of SDSF, in The Forest of Nisene Marks State Park. Numerous cracks and fissures dating from the 1989 earthquake have been located in the State Forest.

Geologic activity, coupled with past fires and severe rain storms, has helped form the steep terrain found throughout the Forest. These events have also contributed to the many landslides present within the inner gorges of streams and along steep roadcuts. The numerous natural springs and sag ponds found throughout the Forest are also the result of past geologic activity.

Table 1. Soil types and associated erosion hazard of SDSF.*

SOIL TYPE	PERCENT SLOPE	PERCENT ACRES	EROSION HAZARD
Ben Lomond sandy loam	15-50	1.77	moderate to high
Ben Lomond sandy loam	50-75	11.61	very high
Ben Lomond-Felton Complex	30-50	0.66	high
Ben Lomond-Felton Complex	50-75	15.81	very high
Hecker gravelly loam	50-75	0.48	very high
Lompico-Felton Complex	5-30	7.44	moderate to high
Lompico-Felton Complex	30-50	23.95	high
Lompico-Felton Complex	50-75	14.71	very high
Madonna loam	15-30	0.88	high
Maymen stony loam	15-30	0.06	high
Maymen stony loam	30-75	1.97	high to very high
Nisene-Aptos Complex	15-30	1.90	moderate to high
Nisene-Aptos Complex	30-50	6.76	high
Nisene-Aptos Complex	50-75	7.62	very high
Riverwash	-	2.94	-
Zayante coarse sand	30-50	1.44	moderate to high

*From Soil Survey of Santa Cruz County, California (US Department of Agriculture, 1980)

The elevation of SDSF ranges from 500 feet at the East Branch of Soquel Creek to 2,500 feet at the southeast corner on Santa Rosalia Ridge. The higher elevations occur in the southeast portion of the Forest and decrease along the ridge going southwest.

WATER RESOURCES

The East Branch of Soquel Creek is a perennial stream that flows through the entire length of the Forest. It is fed by the perennial streams of Fern Gulch and Amaya Creeks from the north, and numerous unnamed intermittent and ephemeral streams. The total size of the East Branch watershed is approximately 19 square miles or 12,240 acres.

As previously mentioned, natural springs and sag ponds can be found in the Forest. The two largest springs are Sulphur Springs, located on the Sulphur Springs Trail, and Badger Springs, located near the main picnic area. Badger Springs was at one time a developed water source as is evident by the remains of a spring box and steel pipes scattered around the area. A third spring, located east of Sulphur Springs along Hihn's Mill Road, was created by the 1989 earthquake.

Amaya Pond, a seasonal body of water, is located in the northwestern arm of the Forest. Approximately one-half acre in size, it is located on the east side of Amaya Road, approximately one-third of the way down from Comstock Mill Road. (See Preliminary Biological Assessment of Soquel Demonstration State Forest, Santa Cruz County, California, [Holland et al., 1992] for more details about Amaya Pond.)

The portion of the East Branch that runs through the Forest is well known for its steelhead rearing habitat. The California Department of Fish and Game prohibits angling in this part of the Soquel Creek watershed in order to protect this important resource. The Fisheries chapter of this report contains more information on the creek and its fisheries assets.

Soquel Creek, including the East Branch, is also part of the domestic water supply for the local community. The lower portion of Soquel Creek serves as part of the natural groundwater recharge system for residents' wells and supplies surface water to a number of intakes along the creek. (See the Fisheries and Watershed chapters for additional information on watershed condition, use, and management.)

ZONING AND GENERAL PLAN

SDSF is entirely classified as Timber Production Zone (TPZ). As defined in Government Code Section 51104(g) and consistent with Sections 51112 and 51113, TPZ land is devoted to and used for growing and harvesting timber and other compatible uses as defined in Section 51104(h). Compatible uses include but are not limited to watershed management, fish and wildlife habitat management, and outdoor education and recreation activities.

The Santa Cruz County General Plan designates the Forest land as Mountain Residential. Objectives of this use include protection of natural resources, retention of rural character, and maintenance of a healthy environment.

CHARACTERISTIC FEATURES

A distinctive feature of SDSF is its proximity to the large urban areas of San Francisco Bay,

Monterey Bay, and San Jose. This provides prime opportunities for urban children to experience forestry education on a first-hand basis.

The natural springs of Sulphur and Badger, as well as other unnamed springs, are also special to SDSF. These springs, plus Amaya Pond, enhance particular biotic communities and offer various research opportunities.

The presence of steelhead trout and a portion of the Soquel Creek watershed also contribute to the special characteristics of SDSF. The watershed, second only in size to that of the San Lorenzo River in Santa Cruz county, represents a system limited to the coastal side of the Santa Cruz Mountains. Steelhead trout, an anadromous fish species, are present. Once abundant along the entire west coast, steelhead populations have declined due to habitat loss and several other factors. The East Branch of Soquel Creek, the portion of Soquel Creek that flows through the Forest, supports a steelhead population and its required habitat. A very limited number of coho salmon and their habitat are also present in the watershed.

As mentioned above, the San Andreas Fault and Rift Zone are directly associated with SDSF. The effects of both ancient and contemporary seismic activity are apparent throughout the Forest. The history and future of this very active system make for an interesting addition to SDSF's abundant natural features.

Finally, the Forest contains archaeological and historical sites discovered during on-going archaeological surveys (Dillon, 1992). The Archaeology Chapter of this plan describes the sites and their significance in detail. Both prehistoric and historic, these sites will enhance SDSF's demonstration and education programs.

ADMINISTRATION

THE LEASE

On March 7, 1988, State Controller Gray Davis and the Bank of America settled a thirteen-year long lawsuit over unclaimed bank accounts. The settlement included \$35.7 million in cash and four undeveloped natural parcels in Tehama and Sonoma Counties. The property that is now SDSF was acquired during the settlement process and added to the package.

The settlement properties are held in a trust with the State as the beneficial owner and the Exchange Bank as trustee. The properties can be sold to pay unclaimed funds if they exceed the \$35.7 million in cash set aside for this purpose. It is doubtful, however, that this will ever happen.

At the time of the settlement, The Nature Conservancy (TNC) volunteered to act as steward for these properties. A 25-year lease was developed which stated that TNC would manage these properties and that past land use practices could continue. Any revenues generated from these activities were to pay for property taxes, operations and maintenance, natural resource enhancement, and access improvement projects.

The Nature Conservancy transferred their lease of the Santa Cruz county property (now SDSF) to CAL FIRE on April 18, 1990. CAL FIRE assumed management at that time and a dedication ceremony for SDSF was held on July 13, 1990. In 2013, at the end of the 25-year lease, the property will be transferred permanently to the State, free and clear.

Under the terms of the lease, both the trustees and the Controller have certain rights and responsibilities. The trustee's primary responsibility is to monitor the lessee's performance as managers of the properties. The Controller is responsible for the sale of any or all the properties in the event that cash assets are insufficient to satisfy all claims. As previously mentioned, this is unlikely to ever happen.

CAL FIRE ADMINISTRATION

Authority to *administer and operate* state forests in California comes from the Legislature and is contained in the Public Resources Code (Sections 4631-4664 and 4701-4703). Rules and regulations governing *use* of state forests are contained in the California Code of Regulations (Title 14, Sections 1400-1439 and 1510-1521). The State Board of Forestry and Fire Protection gives policy direction to the Demonstration State Forest Program, which is administered by the Director of CAL FIRE.

CAL FIRE is administratively broken into two Regions, each with a Region Chief who reports to CAL FIRE's Director. Each region includes units, and state forests are administered by a local Unit Chief. SDSF is in the Northern California Region (with headquarters in Redding) and is within the San Mateo-Santa Cruz Unit (headquartered in Felton). The State Forest office is located next to the CAL FIRE Soquel Forest Fire Station at 4750 Soquel-San Jose Road in Soquel, California. When fully staffed, SDSF has a staff of five : Forest Manager, Assistant

Forest Manager, half-time Office Technician , and two seasonal Forestry Aides. The Forest Manager is supervised by the Unit Chief.

The Forest staff is responsible for the on-site operation of the Forest as provided for by the Public Resources Codes, California Code of Regulations, and the State Board of Forestry and Fire Protection. Forest regulations, policy, and other issues prescribed by the Director of CAL FIRE are used to develop plans and procedures to govern development and perform maintenance of the Forest. The General Forest Management Plan will be reviewed and approved by SDSF'S Advisory Committee (described below) and ultimately approved by the Director of CAL FIRE and the Board .

FUNDING AND TAXES

The Demonstration State Forest Program, including SDSF, is funded through the regular annual state Budget Act. AB 1965 did not establish a separate fund for SDSF and it is not listed as a line item in the state budget. Revenues from all state forests are deposited in a special fund called the Forest Resources Improvement Fund (FRIF), providing money for the annual budgets of state forests.

Expenditures for all state forests are included in a single budget line item in the Department's annual budget. Soquel State Forest was added to the Department's state forest budget in the 1990-91 fiscal year with a minimum of staffing and operating expense. The Budget Change Proposal recognized that there would be little revenue from SDSF for the first few years and that FRIF would need to contribute over one million dollars in operating expense before the Forest could produce revenue. It was also recognized that many years would pass before revenues would equal expenses.

The Department may permit a limited amount of commercial timber operations on SDSF in order to provide funds on a cumulative basis as necessary for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF. The enabling legislation requires a minimum level of timber harvesting, or “floor”, to provide income for all costs of operation and for research and educational purposes of SDSF. The legislation authorizes a higher level of harvesting, or “ceiling”, which shall not exceed long-term sustained yield (LTSY) and on a cumulative basis shall not exceed the level of timber harvesting necessary to provide the funds needed for the maintenance and operation expenses, reasonable capital costs, and other expenses incurred in fulfilling all the objectives identified in PRC Sections 4660-4664 on SDSF. These additional objectives include watershed protection and monitoring, and demonstrations of compatible rural land uses and historic development of timbering and forestry machinery.

As a practical matter, the various objectives overlap and cannot be completely separated. For example, demonstration or experimental timber harvesting could qualify as research and public education as well as an example of compatible rural land uses. The protection of old-growth redwood trees will occur under normal operations of SDSF and does not need to be identified as a separate purpose with separate funding.

Funding for SDSF needs to be increased over time in order to fulfill the objectives of PRC 4660-

4664. In order to adhere to the administrative and budgeting processes currently in place while simultaneously assuring compliance with the limitations placed on the Department by the enabling legislation, the Department will publish accurate annual reports which will compile revenues, expenses itemized by program, and cumulative balances. The Department will maintain a mailing list for requests of annual distributions of the report. Large capital expenses (e.g., for additional properties to provide proper access to SDSF or for construction of a Forestry Education Center and administrative facility) will be funded through the FRIF fund or any fund source approved by the Legislature. Amortization of these capital outlays may be included in the computation of cumulative expenses in the annual reports.

The State pays property taxes to the County of Santa Cruz on land values within SDSF. Additionally, purchasers of state forest timber are liable for payment of timber yield taxes according to Public Resources Code, Section 4654. SDSF's timber sale purchasers are required to file quarterly tax returns with the California Board of Equalization.

ADVISORY COMMITTEE

The legislation which created SDSF (AB 1965) called for the establishment of an advisory committee to assist with the development of SDSF's General Forest Management Plan. A main function of the committee is to act as a critical link between CAL FIRE and the community in the planning effort. This allows SDSF to learn what the community expects and how they feel about certain issues.

The original Advisory Committee had nine members appointed by the Director of CAL FIRE in August of 1991. Five positions were specified by the legislation and four were added by the Director. The original committee consisted of one representative from each of the following:

- * State Board of Forestry and Fire Protection
- * Santa Cruz County Board of Supervisors
- * California Department of Parks and Recreation
- * California Department of Fish and Game
- * The Forest of Nisene Marks State Park Citizens Advisory Committee
- * The Nature Conservancy
- * Soquel Creek Water District
- * Neighborhood Representative
- * Local Registered Professional Forester

The Committee held monthly meetings during the development of the 1998 General Forest Management Plan. The Advisory Committee became inactive, but will be reconvened to provide input on updating the Plan. Most of the groups listed above will continue to be represented on the Committee. However, the Soquel Creek Water District removed itself from the Committee and the State Park Citizens Advisory Committee has been disbanded. The latter groups will be replaced with representatives from the Stewards of Soquel Forest and from the Resource Conservation District of Santa Cruz County. Following its approval, the group will meet twice a year or as needed to review progress on plan implementation and contribute to on-going planning activities. Each member serves a three-year term or until the General Forest Management Plan

updating process is completed (whichever is longer). Once these conditions have been met, one third of the members' terms shall expire on the last day of each year.

SAFETY

The remote and rustic character of SDSF makes safety an important management consideration. Forest visitors need to be informed of safety issues and hazards inherent to the Forest. Roads, trails, and facilities are maintained in safe condition. The staff coordinates with local CAL FIRE forest fire stations and the county sheriff's office for emergency medical response; Forest personnel, including volunteers, will be trained in first-aid. Additionally, search and rescue organizations are allowed to train in the Forest so as to develop their skills and better acquaint themselves with the terrain.

The following safety protocols are currently in place:

- * Restriction and regulation signs are posted at Forest entrances. Hazards, safety issues, and the primitive nature of the area are stated on signboards and in the SDSF brochure.
- * The staff works with the California Department of Parks and Recreation to provide trail maintenance, safety, and coordinated emergency response along the common boundary.
- * All trails and roads are regularly inspected. Fallen trees and other hazards are removed as needed to maintain safe conditions.
- * Motorized vehicles are prohibited beyond designated parking areas. Exceptions are made through special permission and for management, patrol, and emergency purposes.
- Coordination with CAL FIRE's Emergency Command Center in Felton and the county sheriff's office ensures effective emergency response in the Forest. All responses for emergency assistance will be recorded and compiled, including calls for police, fire, medical, or search and rescue services. (Refer to Appendix C, Monitoring Plan.)
- * Volunteers from the Stewards of Soquel Forest and the SDSF chapter of the International Mountain Bike Association provide trail maintenance and patrols, assistance to forest visitors, and support for emergency response.

LAW ENFORCEMENT

Forest regulations and policies are posted on signs and enforced through patrol and apprehension of violators (also see the Resource Protection chapter). CAL FIRE peace officers, authorized under the California Penal Code, will be used to detain violators, with local law enforcement agencies providing backup when necessary. The Department of Fish And Game wardens will

enforce fishing, hunting, and trapping laws. Seizure and prosecution of violators shall be actively pursued. Violators will generally be cited and expected to appear in court, but may be taken into custody if warranted. Methods to prevent illegal activities and alternatives which curtail unwanted behavior will be explored and developed to reduce law enforcement problems.

State Forest trespass violators will normally be cited under sections of the California Code of Regulation (Title 14). Illegal trespass includes removal of trees without a permit and parking, camping/campers, or building in the Forest.

Marijuana cultivation in the Forest has been relatively minor. Since the dedication of the Forest, the remnants of nine old gardens have been found and four active gardens have been eradicated. The general inactivity of marijuana cultivation is due in part to the majority of the Forest being south of the East Branch of Soquel Creek with a northern exposure. Furthermore, SDSF is inaccessible by vehicles and, therefore, is not readily available for use.

Detection of marijuana gardens will occur during normal patrol activities or as leads are developed. Most detection efforts will depend on flights by the Santa Cruz County Sheriff's Marijuana Eradication Program. Information about gardens found in SDSF will be referred to the sheriff's office.

ACCESS NEEDS

As stated in the Soquel Demonstration State Forest Recreation Study Final Report (McNally and Hester, 1993) and the Recreation chapter, Forest access is a significant problem and complex issue. The inability of the public to drive to the forest when county roads are closed and trespass across private property to and from the Forest comprises the bulk of these problems.

The SDSF property came into state management with two verified legal access routes (see the Roads and Other Improvements chapter for more details). The first is an administrative and public right-of-way through the Burch property off Highland Way. The second is across CHY Company property, through the Olive Springs Quarry, which includes a right-of-way for administrative use only.

Many visitors come in via Ridge Trail from The Forest of Nisene Marks State Park. Some recreationists, particularly equestrians, have permission to pass through private property to enter the Forest. The public's use of Comstock Mill Road is prohibited because road conditions are unfavorable.

There is an obvious need to develop better access into the State Forest, particularly along the south and west ends where most trespassing occurs. It is important to provide Forest visitors with additional safe and legal access in order to reduce trespass onto private property. Furthermore, alternative access points are necessary when landslides close Highland Way and/or Eureka Canyon Road..

LAND ACQUISITION PRIORITIES

Land acquisition which would improve access to SDSF is a top priority. This has been actively pursued since the dedication of the Forest. Negotiations with various neighboring property owners regarding appropriation and/or easements have been ongoing. Parcels formerly owned by the Noren family were purchased in 2006 and purchasing land from CHY Company will be considered. Any land acquisitions funded by timber harvest revenues from SDSF shall be limited to those which improve access to SDSF or otherwise directly enhance SDSF. All acquisitions for SDSF will comply with CEQA.

Another acquisition priority is the Burch property which contains the entrance to the Forest (see Figure 2 in the Property Description chapter). The benefits of this procurement would include access control, additional areas for resource management and recreational visitor use, and a location for an informational kiosk and restroom. There would also be clear authority and responsibility for maintenance of and improvements to the bridge, roads, and parking area.

A third priority is the purchase of an area for SDSF's Forestry Education Center (FEC; see the Demonstration and Education chapter for more information). Further study is needed, however, to determine the overall function of the FEC and the facilities and programs necessary to render it successful. Once this has been completed, the amount of property needed and the best site can be determined.

COOPERATION WITH ADJACENT LANDOWNERS

As the previous CAL FIRE Director, Harold Walt, indicated at SDSF's dedication ceremony in 1990, it is important for SDSF to cooperate with their neighbors. Forest staff will continually work with the community regarding local issues. These issues include fire prevention, trespass, watershed impacts, fisheries restoration in the East Branch of Soquel Creek, and mitigation of recreational and timber harvest impacts. The staff already works with local schools in forestry education and also coordinates fire prevention programs with the local CAL FIRE Unit.

COOPERATION WITH OTHER AGENCIES AND INSTITUTIONS

SDSF cooperates with other agencies in resource protection, fire prevention and suppression, law enforcement, and safety. Cooperation is also encouraged for demonstration, forestry education, and university research projects. Other agencies that work with SDSF include the California Department of Parks and Recreation, California Department of Fish and Game, County of Santa Cruz, and the Resource Conservation District of Santa Cruz County. Cooperative institutions of learning and research include UC Santa Cruz, Cabrillo College, California Polytechnic State University at San Luis Obispo, San Jose State University, UC Berkeley, and local public schools.

PUBLIC COMMENTS AND CONCERNS

Public comments and concerns involving the administration of SDSF have largely been focused on the issue of safe and legal access but have also touched upon the FRIF program and Advisory Committee composition.

Suitable public access into SDSF is a major concern for neighbors, users, and Forest staff. As previously stated, there is an obvious need to develop adequate entry and exit points and to stop illegal trespass. Several individuals have made specific requests regarding the location of access points, usually focusing on the area from Olive Springs Quarry to Comstock Mill Road. However, neither end is owned or controlled by the state and only administrative access is allowed. SDSF will continue working with its neighbors and on potential access acquisitions to alleviate this complex and pressing issue.

During the process of establishing SDSF and creating the General Forest Management Plan, questions regarding the purpose and use of FRIF monies have been raised. As required by the Public Resources Code, all revenues from SDSF's timber sales must go into FRIF to be managed and allocated. As part of a state agency, SDSF will adhere to the requirements of FRIF as outlined by law.

Finally, neighbors of the State Forest have expressed that they would like greater representation on SDSF's Advisory Committee. Letters were written to former CAL FIRE Director Richard Wilson by both neighbors and the committee chairman requesting a review of the public's concerns. After a careful and lengthy evaluation, the Director determined that the committee composition was adequate due to strong current local representation.

BIOTA

INTRODUCTION

Biota are defined as the flora (vegetation) and fauna (wildlife) which inhabit a particular area. When people envision forests, such as SDSF, biotic elements are what generally come to mind. Biota, however, are only one element of an ecosystem, small pieces of the larger puzzle.

An ecosystem has been defined as "the interacting populations of plants, animals, and microorganisms occupying an area, plus their physical environment." (Hunter, 1990). The physical environment consists of abiotic factors such as soil, water, space, and climate. The presence and actions of humans make up a yet another component of ecosystems. Though humans often consider themselves to be separate entities, at a basic level they are part of the biota and are intricately connected to everything in an ecosystem. When biotic, abiotic, and human components of a forest ecosystem are working together in dynamic balance, diverse biota and healthy forests are attained. (Dynamic balance refers to the continual interaction of ecosystem components which leads to a balanced yet constant state of change.) In real life, there are rarely clear boundaries between adjacent habitat communities or even ecosystems. Ecosystems and habitats blend and overlap but can be given a label based on general wildlife, vegetation, and location features. Management planning for ecosystems or habitat communities which looks at landscape patterns caused by this overlap can benefit all inhabitants, whether they be stationary (e.g., plants) or mobile (e.g., animals).

In this chapter, the biota are described for each habitat type, or community, found in the Forest. Only the most common plants and animals found in these communities are mentioned, but complete flora and fauna lists are located in Appendix B. For a detailed account of the biota of SDSF, refer to Preliminary Biological Assessment of Soquel Demonstration Forest, Santa Cruz County, California (Holland et al., 1992). For specific information about trees (size, abundance, etc.), see the Timber Management chapter.

Fisheries resources are described in a separate chapter that also deals with aquatic invertebrates. Though terrestrial invertebrates (e.g., terrestrial insects) are critical to terrestrial vertebrate communities, very little inventory or monitoring work has been done to date in SDSF. In the future, more work will be completed in this field. In fact, the State Forest offers prime opportunities for study and research involving terrestrial invertebrates.

HABITAT COMMUNITIES

Communities consist of the living organisms collectively found in an ecosystem (Hunter, 1990). Even though considerable overlap often occurs, communities are individually labeled and classified for research, inventory, and education purposes. They are generally named for the dominant plant species within each community. The dominant plant species is dependent on specific environmental conditions (e.g., soil, climate, water) that further characterize the community. Because communities overlap, plants indicative of one habitat type may be found in others. Poison oak, for example, can be found growing in virtually all of the communities of

SDSF but is most abundant in drier habitat types. Also, some fauna considered to be permanent residents of a particular community actually travel through several communities. These animals, including large mammals (deer, bobcat, gray fox, mountain lion), can be found throughout SDSF as they search for food, water, and shelter.

COAST REDWOOD FOREST COMMUNITY

The coast redwood forest is dependent on areas of high moisture. In its northern range, large continuous stands of redwood are formed. Southern redwood forests are restricted to moist canyon slopes and riparian zones since more moisture is available in these areas. As its name suggests, the dominant tree of this community is coast redwood, the majority of which in the State Forest are second growth (regrowth after original clearcutting). Small groves of old-growth redwoods occur at Badger and Sulphur Springs, and individual old-growth redwood trees are scattered throughout the Forest. Other common trees of this community are tanoak, Douglas-fir, and madrone. Understory species found in the redwood community of SDSF are redwood sorrel, California hazel, wild ginger, and western sword fern. Common wildlife residents include pygmy nuthatch, Steller's jay, and Trowbridge's shrew. The redwood community provides nesting habitat, cover, and food for birds and a variety of small mammals. Redwood forests have comparatively little forage value for deer and other large mammals, however, limiting use of this habitat type.

MIXED EVERGREEN FOREST COMMUNITY

This forest community is dominated by tanoak and Douglas-fir; secondary tree species are madrone and California bay (Holland et al., 1992). Mixed evergreen forests are on the drier slopes above the redwood community, though the two overlap considerably. In fact, the presence or absence of redwood is the only significant difference in dominant vegetation (Holland et al., 1992). Mixed evergreen refers to a combination of conifer and hardwood trees that do not drop their leaves in the fall. Common understory species include poison oak, California blackberry, vetch, toyon, and yerba buena. Familiar wildlife residents are Merriam's chipmunk, dusky-footed woodrat, western gray squirrel, California slender salamander, acorn woodpecker, sharp-shinned hawk, and screech owl. Evidence of feral pig activity is also found throughout the community. Oak trees located in these areas have very high value for wildlife, providing both nesting and roosting substrates and a food source through acorn production.

RIPARIAN COMMUNITY

Riparian communities are named for the intermittent or continual presence of fresh water rather than the vegetation of such areas. Riparian communities are located along the edges and floodplains of streams or surrounding lakes. In SDSF, an abundant riparian community exists along the floodplain of the East Branch of Soquel Creek and to a lesser degree along Amaya Creek. This community is dominated by deciduous hardwoods such as white alder, bigleaf maple, black cottonwood, and California sycamore. Along with these trees, red and yellow willows grow in dense clumps along the banks of the East Branch. Horsetails and hedge nettles are common ground cover along the edges. Wildlife residents include vireos, warblers, Pacific-slope flycatcher, long-tailed weasel, and raccoon. Pacific newts, brown-colored salamanders with bright orange bellies, are abundant in the riparian community and a great delight to young forest visitors. The Pacific tree frog, thought to be common, is only found in a few locations

(Holland et al., 1992). Additionally, large colonies of ladybug beetles gather along creeks to overwinter and breed.

Riparian communities are the most productive terrestrial habitat type for wildlife because of structural diversity and the presence of water. Many migratory songbirds are dependent on riparian habitat for breeding and foraging. Large mammals use the riparian zone as a water supply, and incorporate it into their home ranges. The riparian community is probably the most significant habitat type in the Forest due to its high value to wildlife and limited regional occurrence.

OTHER COMMUNITIES AND ADDITIONAL SPECIES

There are a few other limited communities present in SDSF. Freshwater marshes are areas where the soil stays wet the majority of the year, supporting characteristic vegetation. These marshes usually occur along the perimeter of ponds, at springs, near shallow pools of streams, or in areas of high water tables. The freshwater marsh community in SDSF is scattered, including only Amaya Pond and a few natural springs (Sulphur, Badger, and a couple of small, unnamed springs caused by or increased from the Loma Prieta earthquake). Wildlife residents include migratory waterfowl, great blue heron, black phoebe, belted kingfisher, and garter snakes.

Less than one acre of SDSF are comprised of grassland communities. Some of these areas are natural, due to soil conditions conducive to permanent grassland establishment. Other grassland areas are the result of past disturbance. SDSF's grasslands primarily contain wild oats and annual fescue grasses. Most grasslands are being encroached upon by coyote brush, lupine, poison oak, and Douglas-fir. Common wildlife residents of the grasslands are the gopher snake and Botta pocket gopher.

The chaparral community exists along the exposed ridge tops and on south-facing slopes at higher elevations. These dry locations support the fire-resistant woody shrubs of manzanita, buck brush, coyote brush, and chamise. Common wildlife residents are Bewick's wren, California towhee, scrub jay, western fence lizard, and brush rabbit.

Exotic (i.e., non-native) plant species occur in disturbed areas along roads and the picnic area at Badger Springs. The most common exotics are French broom, periwinkle, jubata grass, and *Eupatorium*. Most exotic species have little value for native wildlife and displace native plant species that have higher intrinsic and wildlife values.

FUNGAL RESOURCES

A local mycological organization has identified a wide variety of mushrooms in the Forest (refer to Appendix B). Fungi are broken into three categories based on their relationship to the immediate environment: mycorrhizal, saprophytic, and parasitic. Saprophytic fungi occur on wood that is already dead whereas parasitic fungi attack and can kill live trees. The most common fungi found in SDSF are mycorrhizal.

Mycorrhizal species form a symbiotic relationship with the trees they grow under. These organisms grow around the rootlets and collect water and trace nutrients for use by trees. The

trees in turn provide carbohydrates to the fungi. Trees and mycorrhizae, therefore, depend on one another for optimum health. According to a representative of the Fungus Federation, , SDSF is a reasonably healthy forest because of the wide occurrence of mycorrhizal fungi (Nathan Wilson, personal communication, 1993).

In general, mycological research of California's wildlands has been minimal, particularly regarding conditions conducive to fungi growth. SDSF provides ample opportunities for mycological research including fungi population analysis and succession of fungal species in the Forest. To protect the fungal resources of SDSF, a permit system for mushroom collection will be maintained and the effects of collection monitored.

PLANT SPECIES OF CONCERN

The Natural Diversity Data Base (NDDDB) was queried on May 10, 2010 to collect information on listed species and species of concern known to occur in the Laurel Quadrangle that contains the Sequoia Demonstration State Forest. A total of eight plant species are state and/or federally listed as threatened or endangered. In addition six plant species are categorized as CNPS (California Native Plant Society) List 1B. The plants of List 1B are rare throughout their range with the majority of them endemic to California. Most of the plants of List 1B have declined significantly over the last century. List 1B plants constitute the majority of the plants in CNPS' Inventory with more than 1,000 plants assigned to this category.

All of the plants constituting List 1B meet the definitions of Sec. 1901, Chapter 10 (Native Plant Protection Act) or Secs. 2062 and 2067 (California Endangered Species Act) of the California Department of Fish and Game Code, and are eligible for state listing. These species must be fully considered during preparation of environmental documents.

A nine quad search of processed CNDDDB data, centered on the Laurel quadrangle identified 48 plant species. One plant species is CNPS List 1A, 32 CNPS List 1B, and 15 are federal and/or state listed as threatened or endangered. While it is unlikely that all or even most of these species would find suitable habitat on Sequoia Demonstration State Forest, the number of species listed provide a rough indicator of extent of plant species of concern in the general vicinity of the State Forest.

SPECIAL-STATUS WILDLIFE SPECIES

The biological assessment of the Forest conducted in 1991-92 found no threatened or endangered plant or wildlife species. Steelhead and coho salmon have subsequently been federally listed as threatened. The California red-legged frog has also been found on the Forest and is now federally listed as threatened. Additionally, a few wildlife species of special concern to the State of California have been seen in SDSF (Holland et al., 1992 and CAL FIRE, 1994). Those species observed were the foothill yellow-legged frog, western pond turtle, sharp-shinned hawk, Cooper's hawk, and golden eagle. The long-eared owl and yellow warbler may also occur in the Forest, but they have not been observed (Table 2). Suitable breeding, nesting, or foraging habitats exist in the Forest for all species observed except the golden eagle.

Table 2. Special-status vertebrate fish and wildlife species occurring or with potential to occur at Soquel Demonstration State Forest.

SPECIES	LEGAL STATUS ^a FEDERAL/STATE ASSOCIATION	HABITAT	OCCURRENCE IN SDSF ^b
Marbled murrelet	T/E	Nests in old-growth conifer forest; forages in pelagic habitats	4
Golden eagle	BCC/FP	Nests in cliffs and trees in forests and woodlands; forages in grasslands, shrublands, and chaparral	2
Cooper's hawk	--/WL	Nests and forages in woodlands and forests; also forages in open habitats	2
Sharp-shinned hawk	--/WL	Nests and forages in conifer forest habitats	1
Long-eared owl	--/CSC	Nests and forages in riparian and woodland habitats	2
Olive-sided flycatcher			2
Purple martin	--/CSC	Nests and forages in woodland and forest habitats	4
Vaux's swift			2
Yellow warbler	--/CSC	Nests and forages in riparian habitats	2
Foothill yellow-legged frog	--/CSC	Occurs in streams with rocky substrate	1
California red-legged frog	T/CSC	Occurs in slow-moving streams, pools and ponds	2
Western pond turtle	--/CSC	Occurs in pools, ponds, and lakes	2
Steelhead (Central CA Coast ESU)	T/--	East Branch Soquel Creek	2
Coho salmon		East Branch Soquel Creek	
Pallid Bat	--/CSC	Buildings, rock outcrops	3
San Francisco dusky-footed woodrat			2

^a Status codes:

Federal

T = threatened.

E = endangered.

P = proposed for listing as threatened or endangered.

BCC = bird of conservation concern USFWS

State

CSC = species of special concern.

T = threatened.

E = endangered.

WL = Watch List

b Occurrence numbers:

- 1 = confirmed nesting/reproduction.
- 2 = observed.
- 3 = not observed.
- 4 = unlikely to occur.

Sources: Jones and Stokes Associates, Inc., 1996
California Department of Fish and Game Special Animals List, July 2009

California Department of Fish and Game State and Federally Listed Endangered & Threatened Animals of California, January 2010

California Department of Fish and Game, California Natural Diversity Database, Laurel Quadrangle, May 10, 2010

The Laurel Quadrangle CNDDDB query for animal species indicates the occurrence of two federally listed endangered insects and two federal or state listed threatened and endangered fish species. In addition, one amphibian is federally listed as threatened and one amphibian, one reptile, and one mammal species are presently California Department of Fish and Game Species of Special Concern. See Table 2 for further information.

PUBLIC COMMENTS AND CONCERNS

A major theme regarding the management of SDSF's biota involves species evaluation and monitoring, particularly for special status species. Comments have largely focused on fauna though concern about edge vegetation and introduced plant species has been expressed. As stated in the Management Guidelines and Planned Actions below, SDSF plans to monitor selected biotic elements of the Forest and evaluate effects of forest management activities on the condition of those resources.

Another primary concern is the restoration of degraded habitats and maintenance of exceptional resource values

in SDSF. Emphasizing this concern are comments regarding management of habitats individually (e.g., manage riparian habitats separately from mixed evergreen habitats) and

management activities which allow interior forest species to thrive. Many commentors feel that development and management of SDSF should be performed in such a way that biotic resources are preserved or improved. The Management Guidelines and Planned Actions deal with these issues as well.

Finally, input has been received regarding the desire for details on exactly how SDSF will manage and maintain its biotic resources. Since this management plan is intended to be general, specific information on the what, how, why, and when of SDSF management activities will be outlined in other formats. More specific management strategies will be developed for different areas and habitats based on anticipated management endeavors and research and monitoring results.

MANAGEMENT GUIDELINES

1. Ongoing monitoring will be performed to detect listed and special status species. Monitoring will include keeping current with state and federal lists as well as conducting periodic floral and faunal surveys. Inventories will emphasize special-status species expected to be present but not yet observed in SDSF. New findings will be added to current species lists. Every consideration will be given to protecting these species and their habitat as required by law and determined by a qualified biologist.
2. Old-growth trees will be protected as outlined in SDSF's authorizing legislation, AB 1965. Areas of old-growth redwood have been located and protection will be provided in all phases of forest management. Additionally, areas have been designated to promote late-succession stands of trees (see the Timber Management chapter for more details).
3. Restore, maintain, or enhance resource values of native habitat communities to promote natural diversity and stability. Measures to achieve this include:
 - * snag recruitment and retention
 - * preservation of appropriate logs and other woody debris
 - * maintenance of natural ponds and springs
 - * protection of riparian zones for use as movement corridors for wildlife
4. Achieve mutual benefit with timber harvesting, demonstration and education, and recreation programs while respecting native biotic elements. Wildlife habitat improvements, such as those mentioned in Management Guideline 3 above, will be considered during the planning and implementation of timber sales, demonstration and education activities, and recreational facilities. Conversely, possible impacts of present or future wildlife and vegetation management projects on other management objectives will be studied prior to project approval and implementation.
5. Control or eradication of exotic plant species will be incorporated into management activities, as appropriate. Ben Lomond Conservation Camp crews, California Conservation Corps members, and volunteers will help with the removal of exotics whenever possible.
6. Control mushroom collection by issuing permits for scientific, educational, and personal use. Mushroom gathering for commercial purposes will be prohibited.

PLANNED ACTIONS

1. Develop a detailed wildlife plan that focuses on the needs of SDSF's biotic community. The plan will provide for habitat and population assessments, management recommendations, and monitoring techniques. Recommendations from the Preliminary Biological Assessment of Soquel Demonstration Forest, Santa Cruz county, California (Holland, et al, 1992) will be incorporated as appropriate along with information from CAL FIRE, California Department of Fish and Game, and local biologists.

2. Work with computer databases for long-range monitoring. The Wildlife Habitat Relationships (WHR) and Natural Diversity Database (NDDDB) reports will be examined and incorporated into the monitoring process.
3. Conduct preharvest surveys to identify active nest sites of all special-status bird species that may occur in the Forest. Those species that may occur in SDSF are Cooper's hawk, sharp-shinned hawk, long-eared owl, and yellow warbler. In addition, preharvest surveys will be conducted for California red-legged frog, foothill yellow-legged frog, and southwestern pond turtle.
4. Continue to evaluate the feral pig situation, and develop management strategies and actions to diminish existing problems.

FISHERIES

INTRODUCTION

Approximately 8 miles of fish-bearing streams flow through SDSF, including the East Branch of Soquel Creek (5.5 miles), Amaya Creek (2 miles), and Fern Gulch Creek (0.5 miles). The East Branch provides valuable steelhead trout spawning and rearing habitat. This, along with their legally-protected status and cultural values, makes the fisheries of SDSF an important resource.

STEELHEAD TROUT AND COHO SALMON

The fishery resources of greatest concern in SDSF are the steelhead trout and coho salmon. Anadromous fish such as steelhead and coho spawn (mate and lay eggs) in freshwater creeks or rivers but spend most of their adult lives in the ocean.

The cycle begins with the development of eggs into young fish in streams where the adults spawned. Once the eggs hatch, young fish develop in the watercourse and gradually make their way to the ocean. Steelhead trout in this area typically spend two years in fresh water although a few may spend additional years inland before migrating. The length of time spent in streams depends on environmental and genetic factors, and some individuals never migrate (Barnhart, 1986). Coho salmon spend one year in freshwater before going to the ocean. Due to the abundance of food, anadromous fish species experience most of their growth once they have reached the ocean. Steelhead and coho along the California coast usually spend two years in salt water, attaining sexual maturity and storing fat for their journey back up streams. As with all anadromous fish, steelhead trout and coho salmon usually return to the stream from which they hatched to mate and lay eggs. Coho die following spawning, but steelhead swim back to the ocean. Steelhead may repeat the cycle and spawn up to four times, but most repeat spawners do so only twice.

Historically, coho salmon spawned in coastal streams from the Bering Sea of the Arctic and the coast of Japan to the Monterey Bay in California. The steelhead's range extended further to the north coast of Baja California. Steelhead and coho populations have been declining throughout their entire range, both in fresh and salt water, for decades due to a number of factors including habitat loss. In California, numbers decrease from north to south, with the southernmost population of steelhead in the Ventura River (Barnhart, 1986), and coho in Soquel Creek. While the Soquel Creek watershed is within the steelhead and coho's range as discussed below, there has been a considerable decline in numbers for both species.

LOCAL FISH POPULATIONS

Anadromous fish populations in Soquel Creek and along most of the Central Coast have declined significantly since the late 1960s. Large numbers of coho salmon were historically present in Soquel Creek, and reports estimate that runs may have included from one to two hundred fish (Runs are groups of fish that swim upstream around the same time of year to spawn.) Coho salmon were planted into the East Branch of Soquel Creek in the 1930s originating from the Brookdale, Big Creek, Prairie Creek and Fort Seward hatcheries (Anderson, 1995).

In 1995, the California Fish and Game Commission listed coho salmon south of San Francisco Bay as endangered. That listing designation was extended from San Francisco Bay to Punta Gorda, Humboldt County in 2005. Coho salmon along the Central California Coast were also listed under the federal Endangered Species Act as endangered in August of 2005 (NMFS, 2010).

No production/mitigation hatcheries (hatcheries that produce fish with the goal of increasing recreational and commercial harvest or for mitigation purposes) for Central California Coast coho salmon currently exist. The two hatchery operations in the Central California Coast Evolutionarily Significant Unit are captive broodstock facilities operated expressly for conservation and recovery purposes with significant oversight by the California Department of Fish and Game and the National Marine Fisheries Service. Coho salmon reared at these two facilities are listed under the Endangered Species Act. In Santa Cruz County the goals of the captive broodstock program include increasing population size, maintaining genetic diversity, and producing sufficient numbers of fish to promote straying into neighboring streams (Sturm, 2009). Coho salmon were thought to be extinct in Soquel Creek in the 1990s and most of the first decade of this century, until a small population of young of the year fish was observed in 2008 near the entrance of Hinckley Creek below SDSF.

Central California Coast steelhead occurs in Soquel Creek in greater numbers. In February, 2006 this species was listed under the federal Endangered Species Act as threatened for coastal basins from the Russian River, south to and including Soquel Creek (CDFG, 2009) and for naturally spawning populations (e.g. not a product of hatchery operations). The listing excludes the Sacramento-San Joaquin River Basin, as well as two artificial propagation programs: the Don Clausen Fish Hatchery, and Kingfisher Flat Hatchery/ Scott Creek (Monterey Bay Salmon and Trout Project) steelhead hatchery programs. Principal hatchery production for the Central California Coast steelhead originates from the Warm Springs Hatchery on the Russian River and the Monterey Bay Salmon and Trout Project on a tributary of Scott Creek. The most recent planting of hatchery-raised steelhead in SDSF (i.e., the East Branch of Soquel Creek) occurred in the 1930s. Steelhead trout from the Brookdale and Big Creek hatcheries were released in the East Branch of Soquel Creek (M. McCaslin pers. comm.). The Monterey Bay Salmon and Trout Project stocked steelhead annually in the main stem of Soquel Creek (downstream of the SDSF) until recently, and in the early 1980s, released fish near the quarry (just south of the SDSF) (M. McCaslin pers. comm.) Estimated run size for the Central California Coast steelhead in Soquel Creek was 500-800 fish in 1982, but declined to less than 100 fish in 1991 (Reavis, 1991) and 50-100 fish in 1994 (Shuman, 1994). The basis for the estimates put forward by Shuman (1994) is, however, considered questionable by Good et al (2005).

Other species of fish within the boundaries of SDSF include the Pacific lamprey and a small resident rainbow trout population above Ashbury Gulch. According to biologist Jerry Smith, additional species likely to be found in the East Branch include prickly sculpin, coast range sculpin, California roach, Sacramento sucker, and northern threespined stickleback.

In order to help protect remaining steelhead trout populations in Soquel Creek, the California Department of Fish and Game prohibits angling in the East Branch of Soquel Creek (Fish and Game South Central District Regulation, Title 14, Section 7.00 [e] [4]). This regulation was adopted by the California Fish and Game Commission in December of 1981 and became effective in March, 1982. In spite of this regulation, poaching of adult steelhead during winter

spawning is prevalent and continues to affect fish populations. To avoid adverse impacts of public use on fish in SDSF, forest staff and the California Department of Fish and Game will conduct ongoing patrols to enforce prohibitions on fishing and fish harassment.

HABITAT AND RESOURCE CONSIDERATIONS

HABITAT ATTRIBUTES

The basic stream attributes for steelhead and coho spawning, rearing, and migration include cool water temperature, high concentrations of dissolved oxygen, adequate water depth, sufficient pool space, and low sediment levels (Barnhart, 1986 and Anderson, 1995). In addition, suitable hiding cover, food supplies, and access to spawning areas can influence the survival of steelhead and coho. All of these habitat conditions need to be considered when working to restore, maintain, or enhance anadromous populations.

Riparian zones are strips of water-loving vegetation and associated organisms that follow the path of watercourses. Essential to healthy aquatic ecosystems, these zones help maintain favorable water quality and provide important food and habitat conditions. Trees along the water's edge shade the water, maintaining cool temperatures for anadromous fish spawning and rearing, as well as maintaining a favorable microclimate for amphibians. Riparian vegetation also stabilizes streambanks and intercepts eroded materials from upslope, minimizing the amount of sediment that enters the stream. Additionally, vegetation adds food and nutrients to the water for use by aquatic invertebrates. Large woody debris falling into the water provides cover for fish, collects and controls the movement of sediment, and creates pools. Consistent with the Anadromous Salmonid Protection Forest Practice Rules approved in 2009, conifers will be planted in riparian zones, in areas where none exist, to promote long-term recruitment of large instream woody debris. Small intertwined pieces of debris, such as branches and twigs, act as collectors of leaf litter and provide more food for invertebrates.

HABITAT LOSS AND DEGRADATION

The loss of habitat, particularly that for rearing, has had a significant impact on the anadromous fish populations of the Soquel Creek drainage. Habitat has declined due to diversion and overdrafting of water by residents, past logging practices, flood control measures, increased development along the creek (including vegetation removal), pollution, and naturally unstable hillslopes. These elements have increased water temperature, added sediment to the creek, altered natural flow patterns, and decreased water levels during the critical summer months and periods of drought. (See the Watershed Assessment chapter for details on SDSF's watercourse conditions.)

Over the years, the combination of increased sediment and lowered water levels have resulted in a considerable loss of pool space throughout the Soquel Creek watershed. This loss of pool space, along with general fresh and saltwater habitat degradation, may have contributed to the nearly complete disappearance of the coho salmon. Coho population densities along the central coast are extremely low and will need vast assistance and human cooperation to become viable again (NMFS, 2010).

An additional habitat problem in the East Branch involves access to spawning areas. A few log jams within SDSF may not be passable by fish. During periods of high water, fish may be able to overcome these barriers, but conclusive evidence of this does not exist. Careful study and evaluation of the steelhead population and barriers may reveal ways to improve steelhead access. Log jams and other barriers will not be removed unless it is evident that such removal will do more good than harm. Furthermore, removal of log jams would be reviewed and approved by the California Department of Fish and Game.

The diversion and overdrafting of water are significant problems in the Soquel Creek drainage, especially along its lower reaches. Near the town of Soquel, a portion of the creek has dried up on various occasions, including the summers of 1991, 1992, and 1994. Some residents along the creek use the water for agricultural as well as domestic needs. There are water allotment requirements, but the requirements of fish were not considered when maximum amounts for residences and businesses were allocated. Because young steelhead move downstream as upper portions of streams dry out in the summer, additional drying downstream limits available space and foraging areas for the entire population.

Greater public awareness and response regarding the aquatic resources of the Soquel drainage are essential for maintaining a healthy anadromous population. Residents need to conserve water, allow the creek to flow its natural path, preserve riparian corridors, and prevent accelerated erosion. While improving the anadromous fish resource within SDSF is a start, it is not enough: Soquel Creek's fisheries require good habitat conditions along every reach of the watercourse and throughout the watershed. All creek users and neighbors should consider the consequences of their actions and how they can help improve the resource.

MONITORING AND ENHANCEMENT

Fish habitat at SDSF will be enhanced through monitoring of fish populations and their habitat, and implementation of habitat enhancement projects in conjunction with ongoing timber operations. This ongoing inventory, which began in 1993, provided information needed to identify goals and objectives for a Fisheries Management Plan. Potential habitat enhancement sites were identified and mapped. Proposed enhancement structures will be reviewed and approved by the California Department of Fish and Game and a qualified hydrologist before installation. To evaluate the success of enhancement activities, regular monitoring will check levels of effectiveness. If any structure is functioning improperly or is adversely affecting aquatic habitat, Forest staff will redesign, repair, or remove the structure, as needed.

AQUATIC INVERTEBRATES

Aquatic invertebrates are organisms with external skeletons that live in water. They include insects (e.g., mayflies), crustaceans (e.g., crayfish), mollusks (e.g., snails), and freshwater earthworms. An important component of aquatic ecosystems, aquatic invertebrates are an essential part of the food web. They typically act as indicator species of fishery habitat quality and water pollution.

When tree litter falls into the water, microorganisms such as bacteria colonize and decompose the material, creating a slime substance. This substance becomes food for invertebrates, which in turn become food for fish. In this way, aquatic invertebrates provide a link in the food chain

between microorganisms and fish, a connection crucial to fish survival.

In fresh water such as the East Branch, juvenile steelhead feed primarily on immature, aquatic stages of insects but will also feed on adult terrestrial insects (Barnhart, 1986). Steelhead prefer the larvae of mayflies, true flies (e.g., gnats), and caddisflies. They are often opportunistic, however, and may feed on any available insect. Adult steelhead typically do not eat during migration and spawning, instead obtaining energy from fat accumulated while living in the ocean.

Because they function as food, aquatic invertebrates, particularly insects, are key indicators of good fish habitat. In most cases, large amounts and diverse species of aquatic invertebrates signify an adequate food supply, increasing the ability of the stream to support larger and healthier populations of fish. Similarly, invertebrates need adequate amounts of leaf litter falling into the stream (therefore, adequate riparian vegetation) and appropriate conditions to support the microorganisms which convert the litter to a usable form. Because of their specific roles and narrow habitat requirements, invertebrates are useful indicators of stream conditions and changes. Through inventory and monitoring the composition of aquatic invertebrate communities, the health of aquatic ecosystems can be carefully evaluated. The first inventory occurred in October, 1995. Periodic monitoring continues on a regular basis.

PUBLIC COMMENTS AND CONCERNS

Comments and concerns regarding the aquatic resources of Soquel Creek were numerous and often fervid. Given the condition of California's CCC coho salmon and steelhead resource and the significance of the Soquel Creek watershed, one would expect a great concern for its future. The comments, concerns, and suggestions received as a result of this concern are represented by two major categories.

The first and probably most important concern involves the maintenance and enhancement of fisheries and other aquatic resources within SDSF. Individuals have stated that, at the very least, SDSF should maintain the resources as they currently exist. Additionally, suggestions for improving the habitat and, therefore, the steelhead fishery touched on in-stream improvements and careful logging methods. SDSF will assuredly work to enhance all wildlife resources and habitats, including those of steelhead.

The second category of concern is really an extension of the first. Comments related that, in order to maintain and enhance SDSF's aquatic resources, inventories and monitoring of these resources must be performed. Specifically, commentators expressed that biologists familiar with the Soquel Creek watershed should set up and maintain a monitoring system which tracks the effects of SDSF's management activities on the resource. Strategies to ensure maintenance and enhancement of aquatic organisms could in turn be developed from the results of inventory and monitoring activities. SDSF plans to conduct the activities mentioned above as well as research and habitat improvement in order to satisfy these concerns (see Planned Actions below).

MANAGEMENT GUIDELINES

1. Protect stream channels, streambanks, and riparian zones during all management activities. Late-succession management areas (see the Timber Management chapter) have been established along all fish-bearing streams and receive specialized management designed to enhance the riparian zone. This will ensure protection of stream integrity, including the channel, bank, and vegetation as well as fisheries resources.
2. Increase the fisheries potential by improving the spawning and rearing conditions of the East Branch of Soquel Creek within SDSF. Methods for improvement will be identified by the assessment discussed above but may encompass actions such as sediment reduction, barrier removal, and pool creation.
3. Demonstrate that other forest management activities are compatible with the maintenance of healthy fisheries populations and habitats. For example, educational displays will be used in recreation areas to inform visitors of the need to practice special care along watercourses. Also, tours of harvested areas could show standard and advanced erosion control techniques utilized to reduce impacts on watercourses.
4. Help residents of the East Branch watershed learn about the components of a healthy watershed and the importance of its maintenance. Educational programs, either formal or informal, will focus on stream health, riparian zones, and the impacts of human use. Once established, the Forestry Education Center (refer to the Demonstration and Education chapter) will likely be the setting for these programs, allowing residents and other interested individuals to learn about the significance of aquatic ecosystems.

PLANNED ACTIONS

1. A fisheries resources assessment was completed and used to develop a draft Fisheries Management Plan (Berlekamp and Sutfin, 1995). The assessment inventoried the condition of the fisheries population as well as habitat status within SDSF. The management plan includes prioritized improvements, monitoring systems, research opportunities, and funding sources. (Refer to Appendix C, Monitoring Plan.) Subsequent fisheries assessments in the watershed have been completed by the Santa Cruz County Resource Conservation District (Alley, D.W. and Associates, 2003) and by the National Marine Fisheries Service.
2. Provide additional aquatic ecosystem research opportunities in the East Branch and its tributaries. Likely subjects for examination include fisheries populations and habitat, aquatic invertebrate communities, and riparian zone studies. Research opportunities for local high school or college students will be emphasized but professional research will not be discouraged.
3. Coordinate with the County of Santa Cruz, the Resource Conservation District, National Marine Fisheries Service, California Department of Fish and Game, and other groups to complete habitat restoration and maintenance projects. Private groups such as the Monterey Bay Salmon and Trout Project or crews from the Ben Lomond Conservation Camp may help with projects such as channel stabilization, hiding cover retention,

riparian vegetation maintenance, pool creation, and barrier removal.

Restoration and enhancement projects for the Forest recommended in the Soquel Creek Watershed Assessment and Enhancement Project Plan (DW Alley, et al, 2003) and the National Marine Fisheries Service in their draft Recovery Plan for the Evolutionarily Significant Unit of Central California Coast Coho Salmon (NMFS, 2010) will be implemented over time as funding, equipment, and/or personnel become available.

4. Continue fish sampling in Amaya Creek and the East Branch of Soquel Creek. Electrofishing will be the primary method but other procedures such as underwater observation may be incorporated. Fish will be evaluated for species, size, health, and location. (Refer to Appendix C, Monitoring Plan.)
5. Conduct aquatic habitat surveys at least once every ten years in accordance with the California Department of Fish and Game methodologies. Monitor ecological condition of aquatic habitats by sampling invertebrates using the protocol developed by the U.S. Environmental Protection Agency in cooperation with the California Department of Fish and Game. (Refer to Appendix C, Monitoring Plan.)

Monitoring work completed on the Forest to date includes annual steelhead trout population surveys from 1993 to 2010 and annual instream temperature monitoring from 1997 to 2010.

6. Fish habitat enhancement structures will be reviewed and approved by the California Department of Fish and Game and a qualified hydrologist before installation. The removal of fish barriers will be reviewed and approved by the California Department of Fish and Game. Where possible, logging equipment will be utilized to assist in barrier removal and enhancement structure development. (Refer to Appendix C, Monitoring Plan.)
7. Conduct ongoing patrols with the assistance of the California Department of Fish and Game to enforce prohibitions on fish poaching and harassment. Incidents will be recorded, compiled, and evaluated annually to determine significant adverse effects on SDSF fisheries. (Refer to Appendix C, Monitoring Plan.)

DEMONSTRATION AND EDUCATION

INTRODUCTION

As stated in the Timber Management chapter, a goal of SDSF is to demonstrate sustained-yield timber production with innovative forest management practices within the context of local community protection, and subject to the limitation of commercial timber harvesting provided in the legislation (PRC 4660-4664). The intent of this goal is to encourage conscientious forestry practices on private lands and demonstrate these practices to the surrounding urban populations. This can lead to improved attitudes toward our natural resources and forestry in general, enhancing responsible stewardship of our forest lands.

Another goal is to show forest visitors that timber management, forestry education, public recreation, and environmental protection are interrelated and compatible. This will be accomplished through combinations of these programs whenever possible. SDSF's location is well suited for the development of forestry education programs because it is close to schools in both the Monterey and south San Francisco Bay areas. This proximity is ideal for groups to learn the importance of forest ecology and management. Similarly, it offers local landowners and the general public an opportunity to view the protection, management, and utilization of renewable natural resources. Field trips would involve public motor vehicle use.

The Demonstration and Education Programs of SDSF will rely on interpretation, volunteer participation, and the creation of an educational center. Each of these elements, as well as the Demonstration and Education Programs themselves, are outlined in the following paragraphs.

DEMONSTRATION PROGRAM

OVERVIEW

The Demonstration Program is a major focus of SDSF for a number of reasons. Most importantly, it provides an opportunity for the general public and educational groups to observe an active working forest. Forestry demonstration can assist the public in understanding forest management and its role in resource conservation and local economics. Additionally, it reminds us of the many wood products we use daily and the importance of keeping forests healthy and productive. Knowledge such as this can help change the negative public image generally associated with forest management practices. Also, the demonstration of various forest activities can potentially benefit small private landowners in the management, protection, and enhancement of their forest lands.

The opportunity for demonstration projects has increased as SDSF has become better established. The staff remain aware of the demonstration potential at the onset of new activities. Demonstration projects may be short term, with several topics addressed each year, or long term, extending over many years.

The primary consideration of the Demonstration Program is to enhance the public's

understanding and awareness of forest management principles and techniques consistent with environmental protection. Throughout the process of establishing projects, a strong emphasis will be placed on environmental protection. The following are demonstration opportunities at SDSF:

Silvicultural Systems	Erosion Control
Tractor/Cable Harvesting Operations	Hardwood Management
Disease and Insect Management	Growth and Yield
Recreation Management	Riparian Management
Reforestation Methods	Cumulative Effects
Fisheries Protection/Enhancement	Fuelwood Management
Road Construction/Maintenance	Prescribed Burning
Old-Growth Redwood Protection	Habitat Enhancement
Watershed Restoration	Fire Protection

PLANNED ACTIONS

1. Demonstrate forest management practices and ecosystem enhancement techniques. To accomplish this, incorporate an identifiable demonstration feature in timber sale planning and implementation, recreation designs and development, or other forest management activities, as appropriate.
2. Develop and implement outreach programs to contact the general public, school groups, and private landowners for demonstrational opportunities. Encourage visits and tours by interested public groups, individuals, schools, and professional organizations.
3. Establish safe and efficient methods of displaying information from demonstration projects and conducting tours of these areas. Displays should be made available and tours held at times that encourage forest user group's attendance and participation (e.g., summer weekends, evenings).

FORESTRY EDUCATION

OVERVIEW

The results of SDSF's Recreation Study (McNally and Hester, 1993) indicate an encouraging level of interest in the Forest's education potential. In order to intensify this interest, SDSF's Forestry Education Program will be informative and fun, providing as many hands-on experiences as possible. The program will be applied in conjunction with demonstration objectives and occur in many different forms. Forestry education will introduce various target groups to progressive forest management practices, resource protection, logging history, forest ecology, and research. This will be accomplished through literature, indoor presentations, and outdoor programs.

Instructional organizations can learn about SDSF's resources by sponsoring classroom presentations or bringing groups to the Forest. School programs may represent a theme, such as Arbor Day or Earth Day, and include activities both at school and SDSF. Classroom and outdoor presentations might also be subject oriented with topics such as watershed management, forest ecology, soil biology, or wildlife found in the Forest.

An educator training program, similar to the one at Elkhorn Slough Reserve, will provide the means for teachers and their students to experience forestry education without relying on Forest staff. A program such as this will offer workshops for educators on different aspects of SDSF and forest resources in general. After completing training, educators may bring their students to the Forest for non-staffed instructional field trips. This removes the time burden from Forest staff members while allowing instructors to absorb new information and reinforce their own training.

Information for the general public and small private landowners will be disseminated through tours, newsletters, brochures, workshops, and seminars. Newsletters and brochures may cover information such as current events, research projects, or compatible rural land uses. Workshops will provide opportunities to learn about forest management techniques and how to solve problems in a group setting. Finally, seminars will focus on specific topics as well as the presentation of ideas, research results, and group discussion.

PLANNED ACTIONS

1. Develop educational information and programs related to forest management and ecosystem processes. Presentations suitable for groups which include people of diverse backgrounds and varying ages as well as group-specific programs will be developed.
2. Determine the best method to distribute information about the forest resources of SDSF to different user groups. Methods to present facts about SDSF may include oral presentations, written information, and interpretive facilities. The Education Study for the Forest was completed in 1997 (Blazej, 1997)
3. Encourage educational organizations, such as local public schools, to use the Forestry Education Program. Stimulate participation by offering quality educational experiences that are both informative and enjoyable. SDSF should establish working relationships

with educators and their students.

4. Develop a teacher training program that will allow educators to lead their own forestry education programs. Offer workshops on topics such as forest ecology, wildlife habitat requirements, and watershed dynamics which apply directly to SDSF. The training procedure should encourage on-going programs which allow students to work with SDSF over a period of time.

INTERPRETATION

OVERVIEW

The public can gather information about SDSF in a variety of ways while visiting the Forest. Interpretive facilities such as self-guided trails, information boards, and hands-on activities offer the Forest visitor an opportunity to learn the many features of SDSF. Since groups who utilize the Forest differ in their use of its many resources, information that applies to and interests many different individuals is essential. Interpretive facilities are a great way for the staff to meet the important goals of public demonstration and education without being relied upon for instruction.

INTERPRETIVE OPPORTUNITIES

Self-guided walking tours incorporate designated stops that are marked and indicated in a guide book. Stops on trails in SDSF might focus on unique sites and vegetation of the Forest or provide activities that facilitate learning. Interpretive trails will work well in SDSF because hiking is a common activity. The Forest provides many possibilities for educational trails, including a number of old logging trails and roads.

Information boards are effective tools that offer educational and operational information. Boards located at parking areas, popular recreation sites, trail heads, and along trails will provide information in the form of displays and brochures. Facts that relate to general forest news are best suited to this type of interpretive facility, and will include subjects like riparian protection, current logging practices, or forest ecology.

Hands-on activities with touchable items and sensory-oriented exercises for adults and children are another form of forestry education. These activities will either stand alone or be coordinated with self-guided and staff-led tours. For example, a touchable tree round on a self-guided trail may help visitors understand how tree age is determined or how human history compares to that of a tree. Forestry tools such as clinometers or diameter tapes could be used to show visitors how trees are measured. Areas containing hands-on activities should be located near popular recreation sites, picnic areas, and parking areas.

Staff-lead tours will travel over expansive areas not covered by self-guided trails and contain stops that are not visibly marked. Guidebooks and hands-on activities may accompany tours, with staff members offering supplemental information.

On the internet – see <http://www.icogitate.com/~tree/SDSF.htm>.

PLANNED ACTIONS

1. Coordinate planning of interpretive facilities with all forest activities, including recreational use, demonstration projects, resource protection, and timber harvesting. Install interpretive facilities near popular recreation sites, parking locations, and areas receiving silvicultural, enhancement, or restoration treatments.
2. Develop an interpretive plan that addresses the SDSF educational audience, types, numbers, and locations of potential interpretive facilities, and the subject matter to be presented. The plan should also include recommendations for facility development as well as monitoring of facility use.

VOLUNTEER PROGRAM

OVERVIEW

A volunteer program will be used to assist the small Forest staff in several ways including efforts to provide a complete and comprehensive forestry interpretive program. Docents are volunteers who have an interest in both the SDSF and natural resources in general and enjoy sharing their experiences with others.

SDSF's docents will be available for public contact and interpretation in both the Forest and educational settings (such as classrooms). In the field, docents will benefit the Forest by providing visitors with information on current events, facilities, vegetation and wildlife, cultural history, and rules and regulations. Ideally, volunteers will have training in first aid and be able to radio for help during emergencies.

Volunteers will also assist in research and monitoring programs, and maintenance and construction projects. The individual skills and talents of SDSF's volunteers, such as knowledge of local flora and fauna, leadership and interpretive skills, and experience working with people, will be utilized to support the Forest. Through their work, volunteers will both benefit the operation of SDSF and acquire a better appreciation of state forests and natural resources in general.

PLANNED ACTIONS

1. Continue to develop a volunteer program which includes docent recruitment. Volunteers will lead tours, patrol the Forest, and assist with education, research, and monitoring programs.
2. Continue to provide training to familiarize volunteers with the history of the state forest system, SDSF's history and objectives, rules and regulations, patrol procedures, and interpretive skills. General training will also include more specific information relating to SDSF's access routes, vegetation, wildlife, and research and monitoring objectives.

TIMBER MANAGEMENT

INTRODUCTION

The main purpose of SDSF's timber management and harvest program is to conduct demonstrations, education, and research in forest management. Sustained-yield management will be demonstrated, with examples of timber harvesting, that ensures proper land stewardship, improved forest health, and protects and preserves SDSF as an intensively managed, multifaceted research forest, consistent with the legislative goals of PRC Sections 4660-4664.

As stated above, a major purpose of the SDSF timber management program is to demonstrate sustained-yield management with examples of timber harvesting. In simple terms, sustained-yield is the yield of commercial wood that an ownership can produce continuously at a given intensity of management consistent with required environmental protection and which is professionally planned to achieve, over time, a balance between growth and harvest. For SDSF, this plan provides for harvest levels well below the level of growth. SDSF is to be used to demonstrate examples of timber harvesting under sustained yield management while also sustaining or improving air, fish and wildlife, water resources, watersheds, aesthetic values, and recreation. Silvicultural methods and harvest techniques that can be applied under California Forest Practice Rules for the southern sub-district of the Coast Forest District will be researched and utilized to demonstrate sustained-yield management with an uneven-aged forest structure.

The harvesting of timber will comply with the discussion found in the Funding and Taxes section of the Administration Chapter. A secondary goal of harvesting timber is to generate revenue to cover maintenance, operation, and other costs of SDSF. This includes funds needed for research, inventory, monitoring, and rehabilitation projects of the various resources in SDSF. Revenues did not meet expenditures, even at the current minimum levels, during the initial nor the second ten-year period of this plan. It is anticipated that revenues and expenditures will converge in approximately the fifth decade of operations. Specific revenue projections are not made because of the inherent uncertainty of timber values and markets. Annual harvest levels may need to be adjusted from time to time to reflect physical conditions in the SDSF, such as catastrophic events. If a significant drop in timber prices occurs, timber harvesting will not exceed the limits discussed in the Harvest Plan section of this chapter. The Department will attempt to maintain basic operation and maintenance services from the FRIF fund in line with other State Forests.

HARVEST HISTORY

The first timber harvesting in SDSF occurred during the late 1800's when the Forest was part of land owned by F. A. Hihn (see the Property Description chapter for a more detailed account of the history of ownership). Hihn logged old-growth redwood along portions of Amaya Creek and his private Sulphur Springs Road. Upon his death in 1913, Hihn's heirs assumed ownership of his lands and continued limited old-growth redwood and tanoak removal.

In 1924, the property was sold to the Monterey Bay Redwood Company (MBRC). The MBRC harvested approximately 100 million board feet of old-growth redwood from 1926 to 1942.

Between 1926 and 1934, logs were yarded to roads using ground lead cable systems powered by steam donkeys. Logs were taken to the millpond, located south of Olive Springs, on trailers towed by tractors. From 1934 to 1942, logs were yarded by tractor and transferred to the mill by log trucks.

The MBRC sold their property to Glenco Forest Products, later known as the CHY Company, in 1961. During their ownership of the SDSF property, CHY performed a small amount of selective timber harvesting between the Sulphur Springs area and the eastern boundary.

In 1979, the Pelican Timber Company purchased a portion of CHY's land, including what is now SDSF, and prepared extensive harvest plans. Pelicans's Timber Harvest Plans were strongly opposed and, after court battles, expired before large amounts of timber were removed. Pelican did actually harvest a small amount, though, prior to state acquisition of the property. Approximately 15 acres of hardwoods in the Sulphur Springs area and 230 acres of second-growth redwood and hardwoods along the west boundary were selectively harvested between 1979 and 1984.

Since the dedication of SDSF in 1990, four timber harvest plans have been approved. The first, Long Ridge timber harvest plan (THP) logged in 1995, was a 64 acre single tree selection harvest and yielded 500,000 board feet. The Amaya Creek THP, harvested in 1998, was also a selection harvest, but included some small (1/4-1/2 acre) groups. This THP harvested 138 acres and yielded 1.5 million board feet. Operations on the Rim THP are active in 2011. The fourth plan, the Fern Gulch THP, has been approved but not yet harvested.

CONIFER VOLUME INVENTORY

There have been four known timber inventories completed for the SDSF property since the turn of the 20th century. The first inventory was completed in 1916 preceding significant harvest of the area. Timber cruisers from San Francisco were hired prior to logging to determine where to locate mills and the best method of transporting logs to those mills. They also judged how much lumber they could harvest from a particular area. The timber cruisers estimated an average volume of 37,300 board feet per acre for the property that is now SDSF, the majority of which was old-growth redwood.

The second inventory was completed in 1978 for Harwood Products, a potential buyer of the property, to ascertain timber growth and volume. Average stand age was approximated to be anywhere from 30 to 90 years old. The average volume per acre for the SDSF portion of the property was estimated at 13,600 board feet, 90 percent of which was redwood.

The third inventory was conducted by SDSF staff in the summer of 1991. The goals of this variable-radius inventory were to establish current volume and basal area by species and to acquire information on species distribution. The average volume for SDSF was estimated to be 28,850 board feet per acre.

The most recent inventory was completed in the spring of 2006. This was a variable-radius plot inventory with plots placed every five acres on a systemic grid. Data measured included tree characteristics, snags, species size and characteristics for all live trees, and unique features such

as goose pens, fire scars, and broken tops with potential wildlife habitat value. Table 3 contains the results for volume and basal area of commercial conifers. The average age of most forest stands was estimated to be 80 years old, based on harvest history.

Table 3: SOQUEL 2006 Forest Resources Inventory Summary

	Redwood	Douglas-fir	Total Conifer	Hardwood	Total
Trees per acre	180	33.0	213	315	529
Basal Area* per acre	178	35	212	116	328
Volume* per acre (Boardfeet for Conifers) (Cubic Feet for HW $\geq 7.0''$)	34,445	7,996	42,441	2,648	NA
Forest Wide Total Volume	95,261,398	22,114,678	117,376,076	7,323,309	NA

* Basal area, the horizontal cross section of trees measured at four-and-one-half feet above ground, is a measure of forest stocking or density. Conifer volume is measured in board feet; a board foot is a unit of measure one foot by one foot by one inch.

GROWTH

A conifer growth study was conducted at Soquel DSF in 1993. This growth study produced per acre values for redwood and Douglas-fir on a forest wide basis. The 1993 survey consisted of visits to 30 of the plots installed in the 1991 inventory to assess growth. Volume growth was estimated based on measurements of tree height and five- and ten-year radial increment growth. The results of this study indicated that forest wide annual conifer growth averaged 972 board feet per acre, with a standard error of 7.2 percent.

The 1991 and 2006 inventories are independent of each other and therefore not directly comparable. An informal comparison of the 1991 inventory and the 2006 inventory however, suggests that current annual growth on the Forest is on the order of 900 to 1,000 board feet per acre per year.

VEGETATION TYPE

A new vegetation type map for Soquel DSF was developed by staff foresters based on their on-the-ground experience. Supporting data used to develop the vegetation map included historical vegetation type maps, a remotely sensed imagery vegetation classification project (Clinton 2009) and the 2006 inventory plot data. The vegetation map uses a standard California Wildlife Habitat

Relationships classification system (Mayer and Laudenslayer, 1988), that includes tree species, structure, and density. Figure 4 shows the Vegetation Type map of the forest.

SITE QUALITY

A timber site quality map for Soquel DSF was developed by staff foresters. The site polygon boundaries were based on their experience and knowledge of the productive potential of different areas on the Forest, guided by site tree data gathered in the 2006 inventory. Additional site tree data was also gathered to fill in any gaps in the site tree coverage. A total of 186 site trees were measured. Figure 5 shows the timber site class map for the Forest.

SILVICULTURE

FOREST DESCRIPTION


Areas in SDSF eligible for harvest include coast redwood and mixed evergreen (Douglas-fir and hardwood) forest communities. Hardwoods, especially tanoak, are a large component of both communities. The normal process in forest succession is for conifers to overtop and suppress hardwoods, as is presently occurring in SDSF. This event will ultimately move the Forest through its current mid-successional stage into a late-successional condition. The 80-year-old forest is reasonably healthy though overcrowded in some areas (too many trees in any one location). SDSF will research and demonstrate ways to improve forest structure and vigor through timber harvest.

The diameter distribution (tree diameter class plotted against the number of trees per acre) of SDSF resembles a traditional uneven-aged forest even though the forest is really a young, even-aged stand (Figure 5). Uneven-aged forests contain many diameter classes and at least three age classes. These forests are typically managed on a size and structure basis. Even-aged forests contain one to two age classes of trees and are managed according to age.

The diameter class distribution of a truly uneven-aged forest forms an inverse J-shaped curve, indicating decreasing numbers of trees as diameter increases. Smaller diameter classes commonly represent younger trees; the number of those smaller trees which mature into larger and, theoretically, older trees then decreases through time, as the curve shape implies. Having a greater number of trees in smaller diameter classes is preferable in order to ensure that an adequate amount of regeneration is present and can keep the forest viable through periods of natural mortality.

As Figure 6 indicates, there are concentrations and deficits of specific diameter classes in the distribution. This is typical of a forest, such as SDSF, that has traditionally been managed on an even-aged basis. Most of the concentrations result from overcrowding, which suppresses growth (e.g., 20-inch diameter class). Those classes with deficits (e.g., 6 -inch diameter class) contain low numbers because the overcrowded areas take up valuable space and nutrients that other size classes could use. Both the overcrowding and deficiency involved with these particular diameter classes can be remedied through a transition to uneven-aged conditions.

Figure 6. Diameter class distribution for SDSF.



SILVICULTURAL TREATMENTS

Silvicultural treatments involve management decisions and actions which direct forest growth, harvest, and regeneration. Between the late 1920's and early 1940's, SDSF was managed on an even-aged basis with clearcutting and natural regeneration. Currently, forests in Santa Cruz County, including SDSF, are managed on an uneven-aged basis as required by the Coast Forest District's southern sub-district rules of the State's Forest Practice Rules. Two ways to establish and maintain uneven-aged conditions are through single tree or small group selection harvest; both of these methods will continue to be demonstrated by SDSF. Although timber harvesting would focus on the removal of conifers, some hardwoods will also be removed to prevent hardwoods from dominating the residual stands.

To create an uneven-aged forest structure, a desired diameter class distribution will be determined and applied to create the favored structure. Silvicultural treatments will focus on harvesting diameter classes that have excess representation. A transition period in which the desired forest size and age structure is obtained will conclude with the forest exhibiting a diameter distribution indicative of a truly uneven-aged forest. Once established, the uneven-aged forest will be maintained through selective harvest in all of the merchantable diameter and age classes and through natural regeneration. Growing space must be provided for young, middle-aged, and mature trees. The preferred level of tree spacing will require a balance between achieving adequate regeneration and providing vigorous growth. Old-growth trees will not be included in uneven-aged management as they will be considered separately from the rest of the Forest (see the following Old-Growth and Late-Successional Management section).

As trees mature, growth rates level off and slowly decline until natural mortality occurs. Reduction of growth at any age, however, can be amplified by conditions such as overcrowding. Selective harvesting can improve stagnant forest conditions and enhance overall health. For example, removal of some trees from overcrowded stands will enable remaining trees to grow faster because of reduced competition for light, water, and nutrients. Sprouts and seedlings can also become established easier for the same reasons, thereby increasing both size and age diversity.

In most cases, forest regeneration will be allowed to occur naturally. This is the best means for SDSF since the dominant tree species, coast redwood, sprouts quickly and vigorously from remaining stumps given sufficient light. In areas that do not regenerate sufficiently (e.g., areas of Douglas-fir, heavy competition, or group openings), supplemental tree planting will take place. Tree planting will also take place for research and demonstration purposes, experimenting with alternative means of forest regeneration. Redwood or Douglas-fir nursery stock will be favored over hardwood seedlings except in areas where hardwood regeneration is insufficient.

SILVICULTURAL PRESCRIPTIONS AND LANDSCAPES

Silvicultural activities such as harvest and regeneration require planning which is typically outlined in silvicultural prescriptions. Silvicultural prescriptions for SDSF will encourage attention to landscape perspectives and patterns that benefit vegetation, wildlife, and other values.

Management from a landscape perspective focuses on landscape patterns and the responses of biotic (e.g., plants, animals) and abiotic (e.g., soil and water) factors to changes in those patterns. A large part of this type of management involves the response of biota to disturbances (such as unnatural openings or burned areas) in the landscape, also known as edge effect. Edge effect, considered to be both beneficial and detrimental, can have impacts on species that are adapted to interior forested habitats. The use of selection methods for harvest, however, can help minimize negative impacts by eliminating sharp contrasts between areas of harvest and adjacent forested areas.

As part of demonstration and research objectives, SDSF will experiment with different sizes of group selection openings as well as feathering the edges of openings. The main purpose of experiments is to determine the extent of edge effect on biota associated with different harvest levels and sizes of group selection openings. Results of demonstration and research projects will be used in the development of future silvicultural prescriptions and activities.

CUTTING CYCLE

Designated forest stands will be selectively harvested approximately every 20 years. The interval between these harvests is known as the cutting cycle. The length of cutting cycles will depend on tree densities, diameter distributions, growth rates, stand objectives and other parameters. Once these items have been analyzed, definitive cutting cycles can be specified. The Option A document prepared for the Forest indicates that cutting cycles will average approximately 20 years.

Traditionally, harvest at the end of a cutting cycle removes the amount of growth that has

occurred since the previous harvest. This may be true for some stands, however, for most stands within SDSF not all of the growth between cutting cycles will be removed (see the Harvest Plan section). By not removing all of the growth, the amount of volume remaining following each stand entry will increase over time.

SILVICULTURAL TREATMENTS AND NON-TIMBER RESOURCES

One of SDSF's demonstration goals is to display that silvicultural treatments are compatible with aesthetic, biological, and recreation values. Ways to display this compatibility may include harvest methods which are aesthetically acceptable both from a distance and close-up, marking treatment boundaries so that they have minimal visual impact from roads or trails, and providing for habitat improvement through silvicultural prescriptions. These and other examples may be elements of demonstration programs directly related to timber harvest.

OLD-GROWTH AND LATE-SUCCESSION MANAGEMENT

OLD GROWTH

Existing old-growth redwood areas in the Forest have been excluded from all future timber harvesting, as mandated by SDSF's authorizing legislation AB 1965. Figure 7 shows that the areas are quite limited (approximately 25 acres) and broken into two to ten acre blocks. (A buffer zone surrounds each block and is included in the total acreage.) Individual old-growth conifer trees will be identified and marked for preservation during timber harvest preparations; individual tree characteristics, listed in Table 4, will be used to identify these old-growth trees in the Forest. (Table 4 compares tree characteristics for old growth against those of young growth.) No individual characteristic, including age, will be used to make a classification. Instead, combinations of these characteristics will be used to determine if trees are classified as old growth or young growth.

LATE SUCCESSION

Late-succession management areas will be administered to promote the development of functional old-growth habitat characteristics through infrequent, low-intensity timber management activities. These characteristics include multi-level canopy structure which provides vertical and horizontal diversity, stand-age diversity, and large trees, snags, down logs, and other woody debris. Recruitment and retention of these characteristics, which provide important habitat elements for fish and wildlife species, will be attained through silvicultural prescriptions. For example, trees with crowns that extend over the stream will not be harvested unless a qualified hydrologist or wildlife biologist recommends removal of such trees to enhance riparian habitat or hydrologic function. These trees may also be removed when recommended by a certified engineering geologist to help stabilize active slides. WHR classifications will be used to assess and maintain the specific wildlife habitat characteristics of these areas in SDSF.

Like areas of old growth, sections in the Forest designated for late-succession management have been identified and mapped (Figure 7). These areas make up 15 percent of the overall forest, and correspond to a width of 300 feet on each side of the East Branch of Soquel Creek, Amaya, and

Fern Creeks within SDSF boundaries (all Class I watercourses). Timber harvesting in these areas would conform to the following guidelines, as well as the Anadromous Salmonid Protection (ASP) Rules approved by the State Board of Forestry and Fire Protection in 2009:

- * At least 75% total shade canopy in multiple layers would be retained.
- * At least 25% shade canopy in trees at least 24 inches DBH would be retained.
- * All woody riparian (i.e., hydrophytic) vegetation would be retained except where riparian function would be enhanced by removing such vegetation.
- * Large snags (i.e., standing dead trees at least 20 inches DBH and 15 feet tall) or live wildlife trees (i.e., trees that support bird nests or have cavities or large limbs that make them valuable for nesting birds) would be recruited (created from existing healthy trees) or retained at an average density of at least five per acre.
- * At selected locations where conifers are lacking, Douglas-firs and redwoods would be planted to promote long-term recruitment of large woody debris in streams.
- * Downed logs at least 24 inches in diameter and 30 feet long would be retained or recruited by felling trees at an average density of at least two per acre, and total coarse, woody debris would be retained at an average density of at least 10 tons per acre.

The ASP rules for Class I watercourses require:

- No harvest occur within the Core Zone or first 30 feet from the watercourse transition line (WTL),
- 80% post harvest overstory canopy for the Inner Zone, or area from 30 to 100 feet from the WTL,
- A post harvest stand that retains the 13 largest conifer trees (live or dead) on each acre that encompasses the Core and Inner Zones.
- Large trees retained are to be those most conducive to recruitment to the watercourse channel.
- Additionally, harvesting is prohibited in channel migration zones and additional protection measures are mandated where there are flood prone area (see CCR 916.9 (f) in the California Forest Practice Rules).

Site disturbance during harvesting operations in late-succession management areas will be kept to a minimum by restricting tractor use and cable-yarding corridors to predesignated trails. Furthermore, all trees to be harvested and all wildlife trees and downed logs to be retained will be marked. A registered geologist and a qualified hydrologist will also be utilized to review operations during timber harvest planning.

Table 4. Individual redwood and Douglas-fir tree characteristics.

<u>YOUNG GROWTH</u>	<u>OLD GROWTH</u>
Abundant branches or knots in the lower 1/3 of the bole (tree trunk).	Lower third of the bole is relatively free of branches or knot indicators.
Branches are small, 3 inches or less in diameter.	Large branches, many larger than 4 inches in diameter.
The tops of the crown are usually pointed and the branches are usually upturned.	Tops of crown are rounded or flat. Branches may be downturned.
Bark is typical of younger trees as described in dendrology books (shallow bark furrows).	Bark is typical of older mature trees as described in dendrology books (deep bark furrows, bark between furrows often plate-like).
Growth is generally fast, as indicated by large annual rings - usually less than 10 annual rings per inch.	Slow current growth as indicated by a long period of narrow annual rings - usually 15 or more annual rings per inch.
Excessive taper in open grown trees.	Very little taper, even in open grown trees.
Diameter at breast height (DBH) generally less than 48 inches.	Diameter at breast height (DBH) generally greater than 40 inches.
Tree age less than 175 years.	Tree age 175 years or greater.

Source: CAL FIRE

Areas along the creeks were chosen for a number of reasons, including protection of sensitive riparian zones, fisheries resources, and existing Watercourse and Lake Protection Zones (WLPZ). Furthermore, these areas provide habitat corridors and buffers between creeks and nearby harvested areas. Corridors are important for wildlife movement and survival, providing a valuable element to the landscape pattern.

HARVEST SYSTEMS

The terrain where logging activities are to occur will be the key factor in determining whether ground skidding, cable logging, or aerial harvest systems are used. The erosion hazard, slope angle, slope stability, and distance to drainages will also be carefully evaluated in the selection of

harvest systems. Additional factors include access, public safety, aesthetics, timing, noise, environmental mitigation, economic conditions, research, and demonstration aspects. The appropriate harvest system for each timber sale will be identified and utilized based on the above considerations. Refinements on the harvest systems used will be made as necessary to accomplish harvesting in an economical, efficient, and environmentally sound manner.

SYSTEMS FOR USE IN SDSF

Ground skidding logging systems, methods which tow logs on the ground to landings (loading areas), consist of tractor, rubber tire skidder, and horse logging. Horse logging, limited to gentle slopes and modest sized trees, can be used to minimize soil disturbance, damage to residual trees in dense stands, or for demonstrative purposes. Tractors and rubber tire skidders are generally used where slopes average less than 35 percent though steeper slopes will be considered on a site-specific basis. (This only occurs on SDSF after evaluations of slope stability and erosion hazard potential and the ability to mitigate any problems to negligible proportions.)

In SDSF, ground skidding equipment will be restricted to the minimum size capable of moving the harvested timber. Skid trail systems (temporary trails used by ground skidding equipment) will be designed so that existing skid trails are utilized where appropriate. All new and rehabilitated trails will be predesignated and placed at a minimum distance of 100 feet apart. A Certified Engineering Geologist (CEG) will participate in the preparation of all timber harvest plans to review the locations of all proposed roads, skid trails, and landings on unstable soils, unstable areas, and areas where the California Forest Practice Rules require that skid trails be designated before the start of timber operations.

Ground lead, high lead, and skyline logging, commonly referred to as cable logging, use cables to partially suspend logs off the ground while transporting them to designated landings. The cables are attached to a machine, called a yarder, which is equipped with multiple winches and a tower for pulling the cables. A wide range of cable systems are available for logging both large and small timber. In SDSF, cable yarding will generally be used on steep slopes, near drainages, or where road construction is difficult.

Aerial harvest systems incorporate the use helicopters and balloons to move logs from one area to another. These systems are beneficial when the erosion hazard or slope instability are high or when access to an area is limited. In SDSF, aerial systems will be considered primarily for demonstrative purposes and where cable yarding is not possible. Helicopter timber drop-off and landing areas will be located at least 0.5 mile from the nearest occupied dwellings.

A preliminary logging plan for SDSF designated proposed tractor and cable yarding areas, existing and planned roads, and old growth areas (Figure 8). Based on this plan, approximately 1,700 acres are suitable for yarding by tractors and 900 acres are suitable for yarding by cable yarders or helicopters.

HARVEST SYSTEMS AND COMPATIBILITY

Harvesting operations will be planned and implemented to minimize conflicts with neighbors and visitors where possible. To accomplish this, road and skid trail construction will be designed with potential users in mind. For example, specific logging trails can be designed to serve as

future recreation trails where appropriate. Certain logging trails may be closed, however, following harvesting activities where public access is inappropriate (e.g., research or sensitive areas, or where trespass across private property would occur). New roads and skid trails will be carefully located to minimize visual impacts. Additionally, the size and duration of area closures and traffic diversions will be evaluated for each timber sale.

HARVEST PLAN

HARVEST LEVEL

Short Term

SDSF's annual harvest level for the next ten years will average between 800,000 and 900,000 board feet (approximately 30 to 35 percent of forest-wide growth and 35 to 40 percent of growth on ground that is available for timber harvesting). This harvest level is consistent with AB 1965 in that it provides for timber management demonstrations and promises an intensively managed research forest. This harvest rate is substantially lower than SDSF's current conifer growth rate of 2.25 million board feet per year and is approximately one percent of the standing conifer inventory. The maximum harvest level allowed regardless of the market price of timber or revenue needs shall be no more than 40 percent of growth or an average of 900,000 board feet per year.

Long Term

The harvest level is projected to gradually increase from 866,000 board feet per year in the first decade to approximately 1.55 million board feet per year in the fourth and subsequent decades. This gradual increase in harvest levels over the first four decades to the equivalent of 50-60 percent of forest-wide growth shall be the maximum percentage of growth harvested in the fifth and subsequent decades regardless of revenue needs. It is anticipated that the funds needed for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF will only be fulfilled once this harvest level is achieved. An analysis of sustained yield production was completed according to the California Forest Practice Rules (CCR 913.11) to determine the effects of this harvest level over time on average stocking level (i.e., timber volume per acre) and corresponding stand growth rates. This analysis concluded that the proposed harvesting schedule is sustainable over the long term.

Sustained yield production is achieved by balancing growth and harvest over time, maintaining a timber inventory capable of producing the long-term sustained yield (LTSY) (i.e., the harvest level projected to be sustainable after 100 years), and having the projected annual harvest level for all future rolling ten-year periods not exceed the LTSY. These results were demonstrated by projecting stocking conditions, growth, and harvest levels for the planning area over a 100-year period based on the best available information and accepted principles of forest management and statistics.

The land area within SDSF was grouped according to its availability for timber harvesting: unavailable (e.g., inoperable areas, old-growth management areas, and unconvertible hardwood areas); limited availability (e.g., late succession management areas, recreation sites, and unstable areas); and fully available. All long-term sustained yield values are based on the Full and

Limited Availability acres only. Unavailable acres are not included in the analysis.

Table 5. Soquel DSF forest land availability.

Timber Capable	Condition Class	Acres	Percent of Ownership
	Full Availability	1,609	60
	Limited Availability	804	30
	Unavailable	268	9.9
	Subtotal	2,677	99.9
Non-Timber Capable	Non-timber	4	0.1
	Ownership Total	2,681	100

Table 6. Decadal harvest, growth and inventory summary of available acres by planning period.

Period	Year	Inventory, MBF	Harvest, MBF	Growth, MBF
1	2010-2020	111,517	8,656	22,495
2	2020-2030	125,356	10,315	23,730
3	2030-2040	138,771	11,824	26,479
4	2040-2050	153,426	14,485	29,057
5	2050-2060	167,998	14,890	30,007
6	2060-2070	183,114	14,960	30,802
7	2070-2080	198,956	15,458	31,454
8	2080-2090	214,952	15,711	32,340
9	2090-2100	231,580	16,039	32,797
10	2100-2110	248,338	15,992	33,357
11	2110-2120	265,703	16,629	33,658

In the first decade, a total of 804 acres of Fully Available land and 402 acres of Limited Availability land are harvested. The remaining Fully and Limited Availability land will be harvested in the second decade. This pattern is repeated in subsequent decades, resulting in a 20-year average cutting cycle.

As shown in Tables 5 and 6 and Figures 9, 10, and 11 the average stocking level on available acres is projected to increase from 46,000 board feet per acre to an average of 110,000 board feet per acre after ten decades. Based on the stocking conditions projected for the tenth decade, the LTSY for SDSF is 3.37 million board feet per year. The LTSY could be increased if more harvesting occurred in the less-available land class. This analysis shows that the proposed harvesting schedule balances harvest and growth, maintains an inventory capable of producing the LTSY, and does not exceed the LTSY in any future period.

Figure 9. Decadal harvest, decadal growth and inventory of available acres by planning period.




Figure 10. Projected annual conifer growth and harvest on SDSF timberlands with full or limited availability.





Figure 11. Projected aggregate conifer stocking on SDSF timberlands with full or limited availability.



As part of the analysis for each succeeding decade, the short and long term harvest levels will be reevaluated based on the best available information at that time. The level of timber harvesting shall not exceed LTSY and on a cumulative basis shall be limited to the level of timber harvesting necessary to provide the funds needed for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF.

HARVEST PROCESS

It is anticipated that approximately every two years, SDSF staff will plan a timber sale, in accordance with environmental laws, which fulfills the harvest objective. Each plan will specify, at a minimum, the area and volume to be harvested, logging and silvicultural methods to be used, and restoration and protection measures necessary to address environmental concerns. All potential harvests will be evaluated by CAL FIRE Forest Practice staff and will be available for public review.

Once a timber sale has been approved by the CAL FIRE Director, a minimum bid will be established for the sale. The timber designated for harvest will then be sold to the highest bidder. The timber purchaser will enter into a contract with the State which specifies all requirements in detail, including the operation and payment schedule, adherence to applicable laws, and any additional improvements or demonstrations. SDSF staff will administer the sale, assuring that the contractor abides by the contract. Furthermore, CAL FIRE Forest Practice Officers from the San Mateo-Santa Cruz Unit and resource professionals from the other Review Team agencies will conduct a Pre-Harvest Inspection and periodically inspect the operation during logging. Additionally, a Waiver from a Waste Discharge Requirement will be obtained from the Central

Coast Regional Water Quality Control Board prior to commencing timber operations.

The harvest process will be administered through the CAL FIRE Class III permit system for major timber sales. Periodically, small timber sales (less than 100,000 board feet) may be offered for demonstration or research purposes. These harvests would require a Class I minor timber sale permit. Any conifers removed in minor sales or lost through catastrophe (earthquake, wind, fire, flood, or pest infestation) must also be applied to the designated harvest level. In the event of a major catastrophe, salvage logging would occur and the annual harvest level would be recalculated.

TIMBER OPERATIONS MONITORING

To avoid adverse impacts on water quality and fisheries resulting from the discharge of sediment to watercourses attributable to timber operations, SDSF staff will monitor all timber operations (including all harvesting areas and new roads, skid trails, and landings) annually for 5-7 years following completion of the operations. Occurrences of substantial surface erosion (e.g., gullies) or mass wasting (e.g., landslides or slumps) resulting from the operations will be identified and described by a Registered Professional Forester. Each substantial gully or landslide will be evaluated to determine its cause and identify stabilization measures that will be most feasible, effective, and cost effective. Such measures will be implemented within 90 days from the date when the subject site is identified, unless due cause for delay is explained and a reasonable alternative schedule for implementation is proposed by the Forest Manager. Appropriate mitigation measures to be incorporated into future timber operations will be described and specified in future timber harvesting plans to avoid a recurrence of the observed erosion or mass wasting events.


TIMBER MARKETS

The local timber market is largely influenced by the proximity of sawmills and economic conditions. The closest large sawmill to SDSF is the Big Creek Lumber Company mill, located 17 miles north of Santa Cruz near Davenport. The next closest sawmills are in Cloverdale, 165 miles north of Santa Cruz on Highway 101 and the Sequoia Forest Industries mill, located in Dinuba, is 150 miles southeast of the Forest.

The timber market has historically undergone fluctuations as a consequence of economic conditions. Figure 12 indicates dramatic changes in redwood stumpage prices during the last 15 years. (State Board of Equalization, 2010).

Stumpage prices reflect the value of logs delivered to the mill less the costs of logging, hauling, and cleanup. In the 1990's, stumpage prices were increasing, but they declined considerably in the last few years. Factors contributing to the rise in the 1990s included a decline in federal timber sales, reduction of available timber elsewhere, increased forest regulations, and protection of threatened and endangered species. The demand for forest products declined recently with the economic recession and decreased markets for lumber related to the housing market. These factors, along with the cyclical nature of the timber market, are expected to continue and influence market activity.

Figure 12. Timber market stumpage prices.




Timber values for SDSF are not expected to be equal to general redwood/Douglas-fir prices. Bidding competition for SDSF sales will generally be lower because of the scarcity of sawmills in the region. Harvesting operations will usually be more expensive because of the amount of rehabilitation and restoration work planned. For example, existing undersized road culverts will be replaced, existing roads will be reshaped and surfaced to reduce erosion and additional fire hazard reduction work will be performed. These revenue reducing activities will be added to the cost of harvesting.

HARDWOOD MANAGEMENT

The 2006 inventory estimated hardwood basal area to be 116 square feet per acre. Figure 13 represents the hardwood basal area by the three most common species and a grouping of lesser occurring species (live oak, willow, alder, and bigleaf maple). As the graph indicates, tanoak is the most common hardwood species present. Hardwoods are a significant component of SDSF and, on average, comprise 40 percent of the basal area.

Hardwood stands will be evaluated during wildlife and timber management planning. The forest type map discussed in the Conifer Volume Inventory section classifies hardwoods as well as conifers in SDSF. Areas that have always been hardwood will remain so and will be examined for wildlife habitat enhancement and research opportunities. Approximately 100 acres of SDSF timberlands that naturally support conifers but, as a result of past logging and other management, currently support pure or predominantly hardwood stands (primarily tanoaks and madrones) will gradually be harvested and reforested where appropriate by planting redwood and Douglas-fir seedlings. These stands are widely distributed throughout the Forest in units one-quarter to one acre in size. A site specific analysis of potential biotic and wildlife habitat impacts will occur

Figure 13. Estimated basal area for SDSF's hardwood tree species.



prior to all hardwood conversion. Any individual openings will not exceed 0.5 acres.

Another aspect of hardwood management is to examine and experiment with alternative uses for hardwoods. The goal is to efficiently utilize more raw material, targeting portable sawmill owners, timber operators, and private property owners in the process. Some items that can be made from hardwood lumber include furniture, paneling, flooring, and decking. Hardwoods can also be utilized as fuelwood and chips. SDSF will research and demonstrate the management and use of hardwoods as a timber resource.

DEMONSTRATION AND RESEARCH PROGRAMS

SDSF's demonstration and research programs will integrate all facets of timber management consistent with PRC Sections 4660-4664 and subject to the limitations of this management plan. Each timber sale will serve as a demonstration but also may contain a research component. Research projects will evaluate the applicability, practicality, and effectiveness of various strategies of forest and watershed management. Timber management demonstrations will include planning and operational projects as well as tours showing the risks and benefits of timber harvesting. Additional information on projects can be found in the Demonstration and Education and Research chapters.

Findings of research and demonstration projects will be available to the general public, small private landowners, researchers, and the forest products industry. Information will be distributed through research publications, direct mailing, libraries, and posting on the internet (see http://www.fire.ca.gov/resource_mgt_stateforests_publications.php).

SDSF can best serve the general public by providing information and experiences which enhance understanding of forest management principles and good land stewardship. The best way to provide learning opportunities is through exposure to forest management activities. Demonstration subjects might include the complexities of land management, trade-offs involved

with timber harvesting, and the extent of wood products utilized by consumers. With this information, the public can make their own informed decisions regarding the effects of forest management on the environment.

Small private landowners can gather useful information and learn to plan more efficiently by viewing land management alternatives. Visiting project sites and observing results is an effective way to understand management techniques and how they might apply to private land. Of most benefit will be practical, directed projects that clearly demonstrate timber management, natural resource enhancement techniques, and solutions to specific problems.

Timber operators and industrial land managers will have the opportunity to view various harvesting and environmental protection techniques. Demonstration and research project ideas will be solicited from this group. Possible projects may include evaluating methods to enhance forest health and regeneration, testing new harvest and management techniques, and solving operational problems.

MINOR FOREST PRODUCTS

The primary minor forest product that will be sold from the Forest is fuelwood. Fuelwood, primarily from hardwood tree species, will be available on an intermittent basis, usually as the result of other management activities. These activities may include road work, fuel break construction or maintenance, by-products of timber sales, and wildlife habitat improvement projects.

Additional forest products may be sold as they become available. These include salvage sawlogs, redwood split salvage, cull logs, roots and stumps, posts, boughs and other greens. As with the hardwood management program, the use of these products promotes utilization of raw materials that might otherwise accumulate and present safety or fire hazards.

PUBLIC COMMENTS AND CONCERNS

Comments and concerns regarding SDSF's timber management program have been very detailed and encompass a variety of subjects. Since it is impossible to address every comment directly, five broad topics of concern which cover most comments have been identified.

The first and probably most important concern to neighbors and other users of SDSF involves disturbance caused by logging activities. Logging truck traffic on mountain roads, reduced access to the Forest, alteration of the forest viewshed, and noise are sources of apprehension for many individuals. As stated in the management guidelines below, SDSF will strive to reduce possible impacts to forest neighbors and users during all forest management activities.

Another concern that has generated many comments deals with the possibility of negative impacts on Forest wildlife. Comments generally focus on the effects of timber harvesting on wildlife habitat, including concern for streams and adjacent riparian areas. SDSF plans to evaluate and monitor the response of various plant and animal species to forest management activities. Results of studies may include strategies to improve adverse conditions, enhance

mediocre areas, or maintain exceptional situations.

The third area of attention deals with revenue from State Forest harvests. Interested individuals have expressed an opinion that SDSF should only harvest enough to cover basic expenses and that revenue should only be used for these expenses only. CAL FIRE recognizes that PRC Sections 4660-4664 limits the level of harvests on a cumulative basis to that necessary to provide the funds needed for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF. CAL FIRE must also comply with California Forest Practice Rules and Board of Forestry and Fire Protection policy. The Department will not attempt to accurately estimate revenues over any specific time period. As revenues increase over time, the Department will request expenditure augmentations through the normal budget process. Final state forest allocations ultimately rest with the Legislature.

Concern for the natural instability of the Soquel Creek watershed and excessive soil loss is also prevalent. Logging activities in the Forest will adhere to California Forest Practice Rules which limit road and skid trail construction. These regulations require site-specific mitigation as necessary to reduce erosion to minimum levels. Additionally, environmentally sound logging practices and experimental research will be used to minimize damage to this sensitive area.

Finally, neighbors of SDSF would like to see local loggers, trucking companies, and mills perform the felling, hauling, and milling of products from the Forest's timber sales. While this may be ideal, the bid process cannot be limited to local businesses. In some cases, the highest bidder will likely be local, keeping the work and revenue in Santa Cruz County .

MANAGEMENT GUIDELINES

1. Maintain and enhance a healthy forest ecosystem. This includes the monitoring of basic resources and requiring management activities that ensure forest vigor. The Forest will be periodically surveyed on an informal basis for general health, with emphasis on disease and insect activity, tree growth and vigor, and soil stability. Other monitoring activities, such as those developed for fisheries and wildlife, will contribute additional information on the health of individual resources within the ecosystem.
2. Protect and monitor the watershed, soil, fisheries, and wildlife resources during all timber harvesting activity. The fundamental goal is to minimize impacts to these resources through planning and mitigation developed on a site specific basis. Protection measures may include selecting low impact harvest methods, avoiding sensitive areas, and conserving or improving resource integrity. Timber harvesting will not occur during the period of winter operations (October 15 through May 1) in accordance with the ASP Forest Practice Rules..
3. Design timber management activities based on landscape perspectives. Components to consider will include horizontal and vertical forest structure, vegetation density, edge effect, corridor size, and biological diversity.

4. Timber sales will have demonstrational value and include experimental and educational aspects whenever possible. This may include pre-harvest and post-harvest activities as well as actual harvest procedures.
5. Consider neighbor and visitor concerns during all timber harvesting activities. SDSF will strive to reduce excessive noise, visual impacts, and transport activity. Logging methods and haul routes that facilitate reduced disturbance will be evaluated on a site-specific basis and used as appropriate.
6. Demonstrate timber management compatibility and integration with recreation. Whenever possible, design timber sales to minimize conflicts with recreational use and improve recreation facilities such as roads and trails. Additionally, safe recreational behavior during logging activities will be encouraged through signs, direction from SDSF staff, and alternative routes.
7. Each timber harvesting operation will include sediment source remediation. High-priority remediation sites will be considered when selecting areas for upcoming harvests. In some cases, remediation at locations other than timber harvest areas could constitute offsite mitigation for the watershed impacts of harvesting.

PLANNED ACTIONS

1. Harvest between 800,000 and 900,000 board feet per year for the next ten years. This is approximately 30-35 percent of forest - wide growth.
2. Protect all old-growth redwood and Douglas-fir trees in the Forest. Promote the development of functional old-growth habitat characteristics in late-succession management areas within 300 feet of the East Branch of Soquel Creek, Amaya Creek, and Fern Gulch. Follow the new Anadromous Salmonid Protection Rules for protection of Class I, II, and III watercourses to enhance riparian functions and help recover state and federally listed fish species.
3. Identify planned harvest areas for the five - year period 2010 to 2015. Evaluate all possible harvesting and silvicultural methods, new road construction needs, and compatibility with other forest uses for each area under consideration. Planning for future harvests will include consideration of potential impacts to smaller watersheds and their future management to allow for well-designed monitoring of potential impacts on water quality.
4. Refine a forest type map which includes tree species, WHR classifications, forest structure, and vegetation density. As resources permit, develop and maintain a Geographic Information System (GIS) database for the Forest which contains this forest type map. The database will eventually include information on soil characteristics, streams, topography, research sites, roads, trails, facilities, and other improvements.
5. Reinventories of the Forest was completed in 2006. This new inventory collected

information by forest type including timber volume, tree age, growth, mortality, stand structure, and WHR characteristics.

6. Monitor all timber operations annually for 5-7 years following completion of the logging operations. Any substantial surface erosion or mass wasting found will be identified and described by a registered professional forester and evaluated to determine its cause. Stabilization measures that will be most feasible and cost effective will be identified and implemented within 90 days. (Refer to Appendix C, Monitoring Plan.)
7. Convert approximately 16 acres of hardwoods back to conifers over the next ten years. Individual openings will not exceed 0.5 acres.

RESEARCH

INTRODUCTION

It is Board of Forestry and Fire Protection's policy for all State Forests to conduct innovative research in forest management. The purpose of such research is to provide resource management information to the general public, small forest landowners, and the forest products industry. Research opportunities in SDSF are numerous, offering small or large, general or specific, and experimental subjects. Individuals conducting research may represent private organizations, including consulting firms and environmental protection groups, or public agencies, such as institutions of learning or resource-based departments.

A plan to identify specific subjects of concern and research priorities for SDSF needs to be formulated so that continuous and long-range studies can commence. Forest staff will investigate possibilities and rely heavily on what the public, CAL FIRE officials, and resource professionals would like to see analyzed.

COMPLETED PROJECTS (TO BE UPDATED)

Several research projects and surveys, summarized below, have been completed for resources within SDSF. Procedures and results of each study are outlined in their respective chapters with the exception of the Geologic Survey, which is covered in the Property Description chapter.

ARCHAEOLOGICAL STUDIES

A complete surface field survey for archaeological and historical sites was conducted in 1991 by Dr. Brian Dillon. Dr. Dillon is a consulting archaeologist affiliated with the California State University at Northridge. Sites found during this study will be protected for cultural, research, and educational purposes. Additional surveys for archeological and historical sites have been completed for timber harvest plans, road and trail construction, and other projects.

BIOLOGICAL ASSESSMENT

An inventory and assessment of SDSF's biota was completed by biologists from California Polytechnic State University, San Luis Obispo, in 1993. The survey resulted in lists of all plants and animals seen, heard, or tracked in the Forest (refer to Appendix B for species lists). The study, which concentrated on special status species, was under the direction of Professors V.L. Holland and Mike Hanson.

A botanical survey of the Fern Gulch area was completed in 2002 and updated in 2010.

GEOLOGIC SURVEYS

An extensive geologic survey of SDSF involved detailed mapping of geologic features and areas damaged during the Loma Prieta earthquake. This analysis was conducted in 1992 by Michael Manson and Julie Sowma-Bawcom of the California Division of Mines and Geology. Mapped information includes geologic characteristics, landslides, stream orders, and areas where mitigation work can be done. Additional geologic surveys have been completed for timber harvest plans, road and trail construction, and other projects.

RECREATION STUDY

A recreation survey to determine forest recreational users' views and use patterns was sent to the neighbors of the Forest and the local mountain community. Copies of the survey were also distributed in the State Forest and in The Forest of Nisene Marks State Park. Results of the study have been a primary source of public input into the forest management planning process. The principal investigators for the study were Marcia McNally and Randy Hester from the University of California, Berkeley. The Draft Education and Recreation Master Plan was completed in 1996 (University of California, 1996).

TIMBER AND GROWTH INVENTORY

A Forest-wide timber inventory and supplemental growth analysis were conducted to determine the current conditions of forest volume and vigor. The timber inventory, completed in 1991 by SDSF staff, surveyed both conifers and hardwoods on a ten percent sample of the Forest. The growth study used a portion of the plots established by the timber inventory but assessed the current growth rates of conifers only.

A new timber inventory was completed in 2006.

WATERSHED ASSESSMENT

A comprehensive cumulative watershed effects analysis for the East Branch of Soquel Creek watershed was completed by CAL FIRE forest hydrologist Pete Cafferata (Cafferata and Poole, 1993). Chris Poole, a student intern from the University of California, Santa Cruz, and Forest staff assisted with data collection. Subsequent studies and surveys have been done by the California Department of Fish and Game, the Resource Conservation District of Santa Cruz County, the National Marine Fisheries Service, and others. Information obtained from these assessments guides future management activities to maintain or enhance watershed and soil integrity.

FISHERIES ASSESSMENTS

Habitat typing for the Forest's five electrofishing stations has been conducted periodically since 1997.

In-Stream Temperature Monitoring has been conducted annually since 1997.

Steelhead Trout Population Surveys have been conducted annually since 1993.

A study on the movement of large woody debris in Soquel Creek was published in 2003 (Lassette and Kondolf, 2003).

FOREST HEALTH

UC Berkeley staff have completed several studies and monitored the status of Sudden Oak Death on the forest since 2001.

RESEARCH PRIORITIES

Subjects for research include all ecosystems in the Forest but are likely to focus on elements in riparian and coast redwood forest habitats. Studies will concentrate on all components of resource management, including protection, management, and public use. Current research priorities are listed below; as forest management techniques evolve, precedence will adjust accordingly.

WATERSHED MANAGEMENT

- * Monitor the effects of forest management activities on the resources of the East Branch of Soquel Creek watershed within SDSF.
- * Investigate stream enhancement and rehabilitation techniques.
- * Control erosion and stream sedimentation caused by logging, road building, and recreational use.
- * Monitor stream discharge and precipitation in small headwater basins.

TIMBER MANAGEMENT AND FOREST HEALTH

- * Investigate optimal spacing requirements for growth and regeneration.
- * Study hardwood management and Sudden Oak Death.
- * Study old-growth redwood forest communities and ways to achieve late-succession stand characteristics over time.
- * Investigate logging techniques which minimize visual, auditory, and environmental impacts.

FISHERIES AND WILDLIFE

- * Assess the current condition of the fisheries resources within SDSF and document long-term trends.
- * Study methods to improve steelhead and coho rearing habitat conditions.
- * Investigate methods to improve wildlife habitat and provide for healthy biodiversity.

RECREATIONAL USE

- * Monitor environmental impacts of visitors to the Forest.
- * Study the reactions and responses of recreational users and neighbors to all forest

management activities.

PUBLIC COMMENTS AND CONCERNS

The general public's wishes regarding research include desire to be kept informed of SDSF's research plans and actions, both presently and in the future. Specifically, curiosity relating to research subjects, objectives, and investigators (and their affiliation) has been prevalent. To satisfy this concern, SDSF plans to announce current research projects through publications such as the Mountain Network News and other local newspapers. Additionally, newsletters containing information on present activities will be posted on signboards throughout the Forest.

Other public comments have suggested that information derived from research studies should be used to formulate forest management policy and actions. Gaining information which will aid in the management of SDSF is essentially the purpose of research projects; the Forest staff intends to put to use relevant information obtained through research.

A final category of public interest involves the availability of study results. Individuals have expressed that research findings be made available as conclusions that may be applicable to their own lands. As stated in the management guidelines below, results will be available through public libraries, natural resource journals, CAL FIRE publications, the CAL FIRE web site, and direct mailing.

MANAGEMENT GUIDELINES

1. Actively design and carry out continuing scientific studies which refine and improve upon existing state of the art forest land management techniques.
2. Coordinate research projects with other State Forests and local, state, and federal public agencies. Additionally, research opportunities have been and will continue to be provided for universities, industry professionals, and private interest groups. Research may be formal or informal, depending on the party conducting the study and available funding.
3. Assure dissemination of research results in a timely and professional manner. Information gained from studies will be made available to local, state, and federal public agencies as well as resource professionals, forest neighbors, and other interested individuals. Reports may be made available through direct mailing, newsletter articles, public libraries, professional natural resource journals, and the CAL FIRE web site.

PLANNED ACTIONS

1. Develop a comprehensive research plan that evaluates research subjects and priorities, identifies proper audiences, and defines techniques to distribute information effectively and in a timely manner. The plan will encourage innovative research in forest management, resource protection, and recreation. An investigation to develop the plan will examine previous research to determine the extent of what has already been completed in the area as well as topics lacking in information.

2. Serve as an outdoor laboratory for CAL FIRE research projects and encourage investigations by other agencies and educational institutions. Use study results to improve forest practices both in the Forest and statewide.

RECREATION

INTRODUCTION

The legislation creating State Forests (Public Resources Code 4631-4664) and Board of Forestry and Fire Protection policy both state that recreation is to be an integral part of the Demonstration State Forest Program. In addition, SDSF's enabling legislation, AB 1965, states that public enjoyment and open access are to be provided. SDSF's recreational management goal is to provide for uses that are compatible and integrated with resource protection, public education, and forest management while following the guidelines of AB 1965 and Board policy.

Prior to state acquisition of SDSF in 1988, the property was privately owned and used primarily by the owners, their guests, neighbors bordering the property, and frequent trespassers. Users consisted of equestrians, motorcycle riders, four-wheel drive enthusiasts, hunters, target shooters, and a limited number of hikers, mountain bikers, and campers.

Since the establishment of SDSF, recreational use of the Forest has changed. Current groups who utilize the forest are (in order of use) mountain bikers, hikers and walkers, and equestrians. The increase in mountain bikers has resulted from an overflow of riders from The Forest of Nisene Marks State Park (TFNMSP) and community awareness of a new riding place. Other recreational users include picnickers, occasional mushroom gathering groups, bird watchers, environmental organizations, a dog search and rescue training organization, and people coming to enjoy the forest environment. Nonmotorized public access is allowed during daylight hours. Camping, campfires, fishing, hunting, organized races, commercial events, and the use of firearms and motor vehicles are generally not permitted. .

Current forest regulations prohibit all recreational motor vehicle use. The primary grounds for this regulation are poor road alignment and maintenance costs. Other reasons include resource protection, safety, and fire prevention. The enforcement of regulations has decreased the number of motorcyclists to a few violators, and four-wheel drive enthusiast use is now almost non-existent.

The draft Education and Recreation Master Plan was completed in 1996. It calls for development of one site "to be used exclusively education oriented group camping"(University of California, 1996). This campground has been developed and is used on a limited bases (Figure 14). The plan also encouraged development of groups to support recreation and education goals and do trail patrols. The Stewards of Soquel Forest and the SDSF Trail Patrollers have since been formed. These dedicated volunteers donate hundreds of hours of labor annually on a range of activities, including trail maintenance and patrol.

THE RECREATION STUDY

To gain a better idea of current and future recreational users, their views, and related issues, SDSF implemented a recreation study which obtained information from the community and

current users. The study was designed to develop recreational and forest management strategies for SDSF. The results were to provide baseline information about existing and potential recreational users that will be utilized as more detailed recreation planning occurs.

The recreation study involved mailing questionnaires to all individuals who share a boundary with the Forest or live nearby. Additionally, different user groups who frequent the Forest were asked to fill out surveys while visiting SDSF. Information about the Recreation Study and the questionnaire were also published in the Mountain Network News, the Summit community's local newspaper. Of the 6,600 individual questionnaires distributed, 800 were returned.

RESULTS

Included here is a partial summary of the findings and recommendations from the study, and what the staff has done to address these issues. For complete details see the Soquel Demonstration State Forest Recreation Study Final Report (McNally and Hester, 1993).

Access

- * **Findings:** Recreational access is currently a significant problem. Many users trespass through private property to enter and leave the Forest. At this time, only one remote, legal access point that provides parking exists. This entry, the main entrance to SDSF, consists of an easement across private property and is subject to winter closures. Furthermore, those who do not use trails (e.g., physically challenged individuals) are limited to using the edge of the Forest only.
- * **Recommendations:** Providing alternative access requires a complete analysis of access needs and an assessment of land available for easement or acquisition. Accommodating non-trail users should be a consideration in future recreation and education planning.
- * **Actions To Date:** Forest patrols have been increased and citations to trespassers have been issued. Publishers of trail guides have been contacted to correct errors and promote an accurate depiction of public access routes. Additionally, loop trails have been developed to reduce trespass. Alternative access routes are being evaluated.

Signs and Maps

- * **Findings:** A number of survey respondents felt that the Forest needs more directional signs and a better map for orientation purposes.
- * **Recommendations:** A map that correctly depicts Forest trails, roads, and access points is a high priority. More clearly visible signs should be placed throughout the Forest, including the entrance at Highland Way.
- * **Actions To Date:** All high-use loop trails as well as the entrance and main road have been signed and the SDSF brochure and map have been redesigned to correct previous shortcomings.

Shooting and Hunting

- * **Findings:** Illegal shooting at Rattlesnake Gulch, private property across Highland Way from the Forest, was a detriment to Forest visitors. Gunfire could be heard throughout the Forest and many users found this unnerving. Additionally, the noise and danger of gunfire reduced the quality of users' forest experiences.
- * **Recommendations:** No recommendations were made to resolve the shooting problem at Rattlesnake because the area was not within state control.
- * **Actions To Date:** The Forest staff worked extensively with the landowner, County Board of Supervisors, Planning and Public Works Departments, and sheriff's office to solve this problem. The area has now been gated and fenced, resolving the problem significantly. The property was acquired by the Mid-Peninsula Open Space District. A trail-head linking this area to the Bay Area Ridge Trail is planned for future construction.

Neighbors' Concerns

- * **Findings:** Some neighbors are concerned that recreation and timber harvesting will disrupt their quality of life in the mountain community. They are worried that too many people will come from the larger metropolitan areas and impact their roads, increase fire danger, and vandalize property. Some individuals have expressed concern regarding excessive noise and their views being destroyed by timber harvesting.
- * **Recommendations:** Every effort should be made to consider the impact of proposed management activities on surrounding neighbors. Work with the community to aid in their understanding of SDSF as public land with certain legislated mandates that include education, public use, and logging. A demonstration that includes visual impact analysis for preparation of timber harvests could be considered.
- * **Actions To Date:** Several public meetings and tours have been conducted to discuss SDSF mandates and hear public concerns. Access issues are being investigated and detailed recreation and timber management planning have been completed. . Motor vehicle use within the Forest is restricted, limiting the types and levels of recreational use.

The Forest of Nisene Marks State Park

- * **Findings:** Sharing a common boundary with TFNMSP is positive and offers prime recreational opportunities for both facilities. Recreationists, particularly mountain bikers and hikers, are starting to use both facilities in one outing. Some people think the Forest is part of TFNMSP. Management staff of both facilities have a good working relationship with one another.
- * **Recommendations:** The relationship between TFNMSP and the Forest has much potential. Thinking of the common boundary as one when problem solving could lead to beneficial solutions for both facilities, extending the State's resources that much further.
- * **Actions To Date:** Managers of both facilities meet regularly to share information and

solve common problems, strengthening their relationship. As a result of this alliance, the new map has been updated to reflect the connections of the Ridge Trail in SDSF with the Aptos Creek Fire Trail of TFNMSP.

Forestry Education

- * **Findings:** There is an encouraging level of interest in the Forest's educational potential. Local teachers are interested in bringing classes out to the Forest and local residents like the idea of educating the general public about the environment.
- * **Recommendations:** Recreational users should be included in the Forestry Education program. Rest stops for different user groups will need to be located so that interpretive materials can be distributed to all recreationists.
- * **Actions To Date:** Outreach programs have been initiated with local schools and natural history organizations. Forest staff members offer verbal educational information to the public, both in the office and out in the Forest. Draft plans for integrating recreation and education have been completed.

Multi-Use Trails

- * **Findings:** Currently, SDSF does not suffer the problems and conflicts experienced on other public lands that allow hiking, horseback riding, and mountain biking on the same trails.
- * **Recommendations:** Successful multi-use trail systems are rare. The State Forest should work to become a model in effective multi-use trail management. The satisfaction of all user groups should be closely monitored.
- * **Actions To Date:** Forest trails are open to all user groups and trail etiquette is stressed to Forest visitors. A Trail Use And Safety Guide has been developed to foster cooperative use and reduce trespass problems.

EXISTING FACILITIES

TRAIL SYSTEM

The trail system of SDSF consists mostly of old logging roads that have been reopened (Figure 14). Currently, many of the trails are not completely contained within the State Forest. Some trails lead from the State Forest to the State Park and others end at private land. Some trails do connect with other trails, however, and can be used as loop trails. Loop trails are necessary to the trail system as they keep visitors inside the Forest and direct them back through the State Park or to Highland Way, instead of onto private property.

Trails are shared by all recreational users, except for one half-mile of Ridge Trail between Corral Trail and TFNMSP. Horses are not allowed on this upper section to keep them from entering the Park, as they are not permitted due to deed restrictions.

Trail Descriptions

Ridge Trail - 3.3 miles

Ridge trail leads from TFNMSP at the southeast corner of the State Forest and follows Santa Rosalia Ridge to Sawpit Trail. It meanders along the common boundary between the State Forest and TFNMSP. This trail is the only other legal public access route to the Forest aside from the main entrance at Highland Way.

Sulphur Springs Trail - 1.5 miles

This trail starts on Hihn's Mill Road and goes up to Ridge Trail. It was constructed in the 1870's to access the Sulphur Springs Resort. The trail passes remnant cold springs, but all of the old resort buildings have been destroyed or removed. This route also serves as an emergency vehicular access to the helipad on Santa Rosalia Ridge.

Corral Trail - 1.7 miles

Corral trail starts at Sulphur Springs Trail and connects with Ridge Trail toward the southeast end of the Forest. It is named for an old corral, less than half way up the trail, that was used by the Sulphur Springs Resort to house their livestock.

Braille Trail – 1.5 miles

The Braille Trail starts on Hihn's Mill road and connects with Ridge Trail. It was originally constructed by mountain bikers at night where they found their way "by Braille". It is now a sanctioned trail, following an agreement that the bike community would stop building illegal trails.

Tractor Trail - 1.5 miles

This trail was constructed in 1934 when logging methods changed from steam donkey to crawler tractor. It was the first logging road developed to access the slopes of Santa Rosalia Ridge. It starts at Hihn's Mill Road and ends at Ridge Trail.

Sawpit Trail - 1.2 miles

This trail is the most recently developed trail in the Forest. It starts at Hihn's Mill Road and connects with Ridge Trail. It is located one-half mile east of the picnic area at Badger Springs. There is a sawpit along the trail that was used before the turn of the century for cutting logs into lumber.

Long Ridge Trail - 1.5 miles

This trail starts on Hihn's Mill Road and travels to Long Ridge Road. The last 0.1 mile is on private property. This trail is primarily used by neighboring equestrians.

OTHER FACILITIES

Picnic Areas

The Badger Springs picnic site was the first such area in the Forest and was established long before the State assumed management. This area is very scenic and, consequently, is a common destination for Forest visitors. Badger Springs is located along Hihn's Mill Road in the alluvial flats of the East Branch of Soquel Creek. The site has several picnic

tables and a few hitching posts for horses. On the opposite side of the road is the Forest's ten-acre old-growth redwood grove, known as Harrison Grove.

Other picnic area locations are the Forest entrance, Sulphur Springs, and on Ridge Trail between Sulphur Springs Trail and Tractor Trail.

Outhouses

There is one portable outhouse in the Forest, provided and maintained by a generous donation from a mountain bike manufacturer. It is located by the eastern property gate on Hihn's Mill Road. .

Sign Boards

There are three information boards in the Forest that provide brochures, maps, and public information for visitors. The boards are located just inside the Highland Gate entrance, at the Badger Springs picnic area, and on Ridge Trail at the southeast edge of the Forest. The Ridge Trail sign board also provides TFNMSP information as it lies on the boundary between the Park and the Forest.

Parking Area

There is one parking area located just outside the Highland Gate entrance. This area is not in SDSF, but the State has a public-use easement with the owners, Roger and Michelle Burch. The parking area has been rocked, and is generally accessible to the public year round.

PUBLIC COMMENTS AND CONCERNS

Many comments and opinions were gathered during the Recreation Study. Appendix A of the study's final report contains all of the results of the questionnaire as well as comments and concerns of users and neighbors. The major public concerns, such as access, are addressed in the Results portion of this chapter's Recreation Study section as well as in the study itself. For detailed comments, refer to [Soquel Demonstration State Forest Recreation Study Final Report](#).

MANAGEMENT GUIDELINES

1. Public safety will be emphasized in all patrol and management planning activities. Violators of Forest regulations will be cited. Prohibited activities include use of motor vehicles within SDSF, camping or fires without a permit, entering closed areas, and fishing (per Department of Fish and Game Regulations). The Forest is also administratively closed to shooting, hunting, organized races, and commercial events.
2. In accordance with Board of Forestry and Fire Protection policy, recreational facilities will be maintained with minimal development, preserving the rustic and informal characteristics of the Forest. Periodic assessments will be made to ensure that facilities meet users' needs while remaining as natural as possible.

3. Regular maintenance will be provided to ensure the upkeep and safe conditions of all existing facilities, including picnic tables, signboards, parking areas, and trails. Periodic inspection and maintenance of recreational facilities will be performed by Forest staff, Ben Lomond Conservation Camp crews, and volunteers, including the Stewards of Sequel Forest and the SDSF Trail Patrollers.
4. Recreation will be coordinated to achieve integration and compatibility with timber management, resource protection, demonstration and education, and the neighboring community. Demonstrations will show how recreation and timber harvesting, two seemingly conflicting management objectives, can be integrated. Interpretive resources will explain the basics of forest management and the need to protect sensitive resources. In addition, visitors will be directed away from private property through circulation patterns (loop trails), use of proper signing (directional, regulation, no trespassing), and keeping forest attractions away from private property boundaries.
5. SDSF will be managed to provide positive experiences for all recreational users. Forestry education will be an integral part of the recreation experience. Proper trail etiquette between mountain bikers, equestrians, and hikers will be encouraged through signs, educational information, and patrol by staff and volunteers.
6. Management will strive to reduce recreational development and recreational use impacts in SDSF's riparian areas. The majority of riparian areas in the Forest are not suited for recreational use and will not be developed. Remaining areas, however, will not be foreclosed to recreational use or forestry education options. Any recreational improvements in riparian areas will be based on site-specific evaluations.

PLANNED ACTIONS

1. Additional recreation planning beyond the scope of this General Forest Management Plan was completed in 1996. The Education and Recreation Plan was developed based on the results of the Recreation Study. The new plan offers solutions to existing conflicts, including access and user circulation patterns. Additionally, alternatives for recreational use such as a rustic group campground were considered and safety concerns and hazard mitigation examined.
2. Recreational development levels that allow the maintenance of SDSF's rustic qualities were determined through the Education and Recreation Plan.
3. Special volunteer programs to enhance recreational opportunities have been established. These programs incorporate docents to help with interpretation and forest education (see the Demonstration and Education chapter) and patrol the Forest via foot, horse, or mountain bike. Facility development and maintenance will be coordinated with demonstration and education programs.
4. Conduct ongoing patrols of riparian areas including those with existing facilities to enforce prohibitions on vandalism and other damage to riparian habitat related to public use. Implement additional restrictions on public use as needed. (Refer to Appendix C,

Monitoring Plan.)

5. Record and compile descriptions of all reported nuisances caused by public users at SDSF or on adjacent ownerships including, but not limited to, trespass, vandalism, littering, and noise. Implement additional restrictions on public use as needed. (Refer to Appendix C, Monitoring Plan.)
6. Compile annual estimates of public use of SDSF in user days using , patroller reports, surveys, trail counters, and other information. (Refer to Appendix C, Monitoring Plan.)

RESOURCE PROTECTION

FIRE PROTECTION

The CAL FIRE San Mateo-Santa Cruz Unit Chief is responsible for fire protection in the State Forest. The Forest Manager, the Unit's Operations Division Chief for Santa Cruz County, and the local CAL FIRE Battalion Chief will work together to ensure an adequate fire protection program is in place for SDSF. In addition, the Forest staff will work with other agencies as needed to provide fire protection for the Forest.

FIRE HISTORY

A 50-year fire history (1929-1979) of the Santa Cruz Mountains was compiled by Jason Greenlee in 1981. The area which is now SDSF was included in the study, and the following fires occurred during that 50-year period:

- 1933 - 240 acres burned in the Amaya Creek drainage.
- 1936 - 54 acres burned along Santa Rosalia Ridge between Tractor Trail and Sawpit Trail.
- 1936 - 118 acres burned in the Hinkley Creek basin, with approximately 25 acres in the Forest, along the ridgeline at the top of Sulphur Springs Trail.
- 1938 - 87 acres burned along the upper portion of Sulphur Springs Trail.
- 1957 - 168 acres burned from Highland Way to Hihn's Mill Road, just east of Ashbury Gulch. Approximately 50 acres were in the Forest.

In addition, an earlier fire in 1922 started in Hinckley Creek and spread to Soquel Creek, burning a total of 7,000 acres.

The exact causes of these fires are not known, but it is believed that logging practices of the past, coupled with high risk machinery and equipment, caused most of them. Phil Mason, a long-time local resident, remembers that the 1933 fire was caused by a steam donkey engine that caught fire, burning the Amaya Creek drainage.

Modern logging methods and equipment are much safer from a fire perspective, and the risk of fire has been greatly reduced. This is evident from the lack of fires in the Forest during the last 30 years. The only known recent fire occurred in 1970 when the High Bridge, at the southwest edge of the Forest, was set on fire by an arsonist.

Recent large fires in the area include the Lexington Fire in July, 1985; the Summit Fire in May, 2008; and the Loma Fire in October, 2009. Though not in the State Forest, the Lexington fire's impact to the Summit mountain community was severe enough to mention here. An arsonist started the fire which burned 13,800 acres in the Los Gatos Creek Drainage (600 acres were in

Santa Cruz County). Forty-five hundred people were evacuated from their homes, including residents of the Summit area. In the end, 44 structures were destroyed, including houses and outbuildings.

The Summit Fire burned 4,270 acres in the upper portion of the Soquel and Corralitos Creek watersheds on May 22 – 23, 2008. Thirty-five residences and 64 outbuildings were destroyed. Portions of the area burned suffered significant damage and mortality to the understory herbaceous and shrub layers, including ground-level vegetation in riparian areas. The majority of the overstory canopy survived intact. Isolated pockets of mature trees, including hardwoods, redwood, and Douglas-firs, were damaged or killed. Knobcone pine and chaparral in the upper portion of the watershed burned with high intensity and suffered significant mortality (State Emergency Assessment Team, 2008).

The Loma Fire burned 485 acres within the Soquel Creek watershed between October 25 and November 2, 2009. One trailer and two outbuildings were destroyed. The fire started on Loma Prieta Ridge and was pushed southwest by strong northeast winds. The fire slowed when it entered the 2008 Summit Fire boundary. Activity also slowed in unburned forested areas, due to higher humidity and higher fuel moisture. Existing roads provided access and control lines to aid the fire-fighting effort. Portions of the area burned suffered significant damage and mortality to the understory herbaceous and shrub layers, particularly at higher elevations. The majority of the overstory canopy survived intact. Isolated pockets of mature trees, including hardwoods, redwoods, and Douglas-firs, were damaged or killed. Knobcone pine and chaparral in the upper portion of the watershed burned with high intensity and suffered significant mortality. The fire burned through a number of swales and watercourses but higher humidities and fuel moisture levels in these areas kept the intensity of the burn relatively low.

PRE-SUPPRESSION

Pre-suppression is defined as fire protection activities performed before fire occurrence to ensure effective fire suppression. Pre-suppression plans discuss site-specific ways to minimize loss and reduce hazard and risk. The current pre-suppression plan for SDSF will be updated by the local CAL FIRE Battalion Chief with assistance from the Forest Manager. The more comprehensive plan will include the definition and assessment of high risk and hazard areas within the watershed boundaries, maps of fire defense improvements, prevention techniques, and an evaluation of available resources. State Forest staff continue to work on these activities.

Fire Defense Improvements

Fire defense improvements will be strategically located to protect forest land and neighboring properties. Improvements in the State Forest include two water tanks (e.g., 10,000 gallons), shaded fuel breaks, and two helipad locations (Figure 14). . In addition, appropriate signing, fire hazard reduction, and adequate access to roads and trails will be added or maintained. Fire hazard and prevention information as well as Forest regulations will be posted on all information boards. The parking area, information boards, and picnic areas will be treated to reduce fire hazards for safety and demonstration purposes. The major roads and trails in the Forest have been opened to provide access for fire protection purposes.

Shaded fuelbreaks protect high value areas such as forest land, historical sites, and neighboring

property. Typically, they are areas 100 to 300 feet wide where vegetation and other forest fuels have been decreased in order to reduce the rate of spread of an advancing fire. All dead trees and ladder fuels (shrubs and lower tree limbs) are removed and the overstory canopy is thinned to a level where shade will still retard the growth of new ground fuels. The understory is modified so that a low-growing ground cover is retained within the fuelbreak to provide fuels to start a backfire. Whenever possible, fuelbreaks should visually merge with the surrounding landscape, conforming to the natural features of the area. Periodic maintenance will be needed to maintain fuelbreak specifications.

Shaded fuelbreaks in SDSF are being constructed, with the help of crews from Ben Lomond Conservation Camp, along ridges and high use roads and trails. This is to provide safe locations for fire control lines and backfiring; ridgelines are commonly used as control points. In addition, shaded fuelbreaks or the modification of fuels around homes will be considered on a site specific basis, as subsequent planning occurs.

Regulations

Restrictions are in effect for hunting and shooting, smoking, and fires in the Forest. Smoking and fires are not permitted anywhere in the Forest, including parking areas. A possible exception to these regulations would be campfires in a proposed permit-only group campground (to be addressed in the Recreation Plan for SDSF). Hunting and shooting are administratively prohibited in the Forest.

The periods of extreme fire danger for SDSF usually occur from July through October though these periods may be extended by severe weather. During these periods, SDSF will follow the Unit's Red Flag Alert Plan. This is consistent with the plan The Forest of Nisene Marks State Park follows during extreme fire danger conditions. The Forest Manager will coordinate with the Unit Operations Division Chief to determine necessary actions to be employed. The steps include increasing patrols of the Forest, posting red flag alert signs, providing more fire prevention information and awareness of current conditions to Forest visitors, and reducing the number of visitors in the Forest by posting the area as closed.

Education

SDSF staff will coordinate with the Unit Fire Prevention staff for educational purposes. Educational information will be used to reduce the number of human-caused fires within the State Forest. Target groups will include neighbors, visitors, school groups, and local organizations. In general, neighboring property owners pose a risk of human-caused wildland fires in the Forest. They will be encouraged to meet with Burrell and Soquel Forest Fire Station personnel for information on CAL FIRE's Fire Prevention Program in order to minimize the risk of wildfire.

Signs will be posted on all information boards alerting Forest visitors to the dangers of fire and ways they can minimize these dangers. Dangerous Fire Area signs will be posted when appropriate. During regular patrols, visitors will be informed of fire hazards and prevention methods.

School groups and local organizations will be referred to Fire Prevention staff or local fire stations for fire safety information and presentations. Groups visiting the Forest for education programs will be informed of fire safety, hazards, and prevention techniques.

Enforcement

Forest patrol is an important part of fire protection and prevention. SDSF staff will coordinate with the Fire Prevention staff for patrol purposes. Patrols will include public contact, fire detection, and patrol of roads and trails during the fire season. CAL FIRE personnel will be utilized for weekend patrols and major holidays, especially during periods of high fire danger. CAL FIRE peace officers will either provide direct supervision or lead these activities. Additional patrols may be conducted by volunteers as deemed appropriate and safe by CAL FIRE.

Fire suppression cost recovery will be pursued for damages resulting from deliberate and negligent acts of Forest users. Active investigations will be used to locate responsible parties and recover maximum legal damages.

SUPPRESSION

Suppression tactics are based on information from and implementation of the pre-suppression plan. SDSF staff can support initial attack fire control personnel by providing local expertise regarding current road conditions, vegetation, and cultural resources. The staff may also evacuate visitors, close the fire area, perform law enforcement tasks, and assist with media information as appropriate.

Detection

Detection strategies include daily patrols, searching for evidence of fires, and CAL FIRE air flights during extreme fire danger periods or after lightning storms. Also, the Unit's Emergency Command Center will check the Automatic Lightning Detection System (ALDS) for possible strikes in the Forest.

Communication

As part of communication, SDSF will maintain an adequate radio system and stay in close contact with local CAL FIRE fire stations (Burrell, Soquel, and Corralitos). Local CAL FIRE fire control personnel will become familiar with the Forest, its road and trail systems, water sources, and landmarks (for use as reference points) and be advised of any changes that occur.

CAL FIRE's resource tracking system, Computer Aided Dispatch (CAD), will be used to dispatch the appropriate personnel and equipment to any fires on SDSF. The State Forest is a defined response area within CAD. The staff will inform the Emergency Command Center of any changes or updates to the CAD database, including information on roads, access points, and fire defense improvements.

POST-SUPPRESSION

Post-suppression activities include the evaluation of pre-suppression information, suppression actions, and fire line suppression repair. Fire suppression repair involves erosion control actions needed to rehabilitate areas directly impacted by suppression activities (e.g. firelines, roads, helipads, etc.). Rehabilitation involves erosion control and other restoration activities not directly caused by fire suppression activities. Unit personnel will evaluate post-suppression activities on an individual fire basis.

To minimize increases in wildfire risks resulting from increased public use in the Forest, the staff will record and compile descriptions of all wildfires occurring at SDSF. This information will be evaluated annually. If an increase in wildfire frequency occurs, appropriate measures will be implemented as needed to reduce wildfire risk.

PRESCRIBED FIRE

Prescribed fire is the controlled use of fire under specific weather and fuel moisture conditions within a predetermined area. Fire, under these conditions, produces the intensity of heat and rate of spread required to accomplish specific management objectives. These objectives could include fire hazard reduction, silvicultural research, and ecosystem enhancement.

A prescribed fire program that involves these objectives will be evaluated for SDSF. A risk and benefit assessment will be the first step in the evaluation. The greatest risk is the loss of fire control and resulting damage that may occur. Benefits include a reduction in fuel load, removal of exotic plant species, and improvements to vegetation and wildlife habitats. A prescribed fire program must consider the mountain community's concerns and address them effectively. An established prescribed fire program provides research opportunities for both resource management and fire behavior purposes.

If a prescribed fire program is established in SDSF, a weather station will be established and monitored, fuel loads and types will be sampled and mapped, and burning prescriptions will be developed for the different vegetation types found in the Forest.

PUBLIC COMMENTS AND CONCERNS

On April 29, 1993, SDSF held a workshop designed to gather information, comments, and concerns from the public. Based on the comments received, there are four major issues concerning fire protection and SDSF.

The first issue involves concern about logging equipment and its potential to cause fires. As stated earlier in this chapter, modern logging methods and equipment are much safer and less likely to generate fires than historic logging practices. Additionally, logging contractors are required by law to develop a fire protection plan for each THP and to follow the state's Forest Practice Rules regarding fire protection. During harvest activities in SDSF, periodic inspections will be performed by both CAL FIRE Forest Practice Officers and Forest staff.

The risk of fire as a result of recreational use is also a concern. Individuals commented on issues such as limiting the number of people allowed in the Forest during periods of high fire danger and the prohibition of fires and camping at all times. Forest policy is to inform users of fire danger and to discourage activities that may result in increased fire risk. Similarly, camping, hunting and shooting, and motor vehicles are prohibited in SDSF though they may be allowed in the future (to be addressed in SDSF's proposed Recreation Plan).

The third issue raised at the workshop concerns the use of prescribed fire in SDSF. Comments about the consequences of both using and refraining from prescribed burning were received. As mentioned above, a risk and benefit analysis will be completed and analyzed before any

decisions are made in this area. Lastly, comments regarding fire defense improvements vary from concern about how improvements may change the character of SDSF to suggestions for road accessibility. Currently, roads in the Forest are kept clear of combustible vegetation by Ben Lomond Conservation Camp crews and Forest staff. Likewise, the condition of roadbeds are examined periodically and maintained as needed. Though fire defense improvements may change the appearance of some areas, their presence is needed in order to allow timely and effective response should a fire occur.

PLANNED ACTIONS

1. Coordinate with the Unit Battalion Chief and other fire control personnel to develop a comprehensive pre-suppression plan for SDSF.
2. Determine fire defense improvement locations and initiate their construction. Include the Unit Operations Officer-South in all decisions.
3. Provide patrols to enforce fire prevention policies, coordinating with the Fire Prevention staff as needed. Forest staff and volunteers will patrol SDSF on weekends and holidays, especially during periods of high fire danger. The Forest will be closed to public use when fire risks become excessive.
4. Conduct fire prevention education for neighbors and Forest visitors utilizing SDSF and fire prevention staff members.
5. Work with the Department of Parks and Recreation personnel from The Forest of Nisene Marks State Park to ensure effective fire protection along Santa Rosalia Ridge.
6. Record and compile descriptions of all wildfires occurring at SDSF including ambient weather and fire hazard conditions, ignition source, area and vegetation types burned, and estimated damage. Implement appropriate measures as needed to reduce wildfire risks. (Refer to Appendix C, Monitoring Plan.)
7. Evaluate the needs and feasibility of a prescribed burn program.

FOREST PEST MANAGEMENT

Forest pests, such as insects, diseases, and vertebrates, have long been established in California's native timberlands. Populations of pests are dynamic and fluctuate in response to climatic and environmental changes such as drought, forest overstocking and windthrow, fire, and other site disturbances. The actual or potential effects of pests may reduce or threaten to reduce anticipated tree growth, species composition, or forest stocking. At the same time, other forest resources, such as wildlife habitat, may be impacted. Integrated forest pest management provides a means to address these issues.

The intent of integrated pest management (IPM) is to prevent or restrain forest pest problems using population suppression and the minimization of factors that predispose trees to infestation. IPM makes use of the benefits of cultural, mechanical, chemical, semiochemical (i.e., synthetic pheromone), and biological pest management alternatives.

Pests known to have caused tree mortality within or adjacent to SDSF are listed in Table 7.

Table 7. Forest pests of the Soquel Creek watershed.

PEST	HOST
Fungal Diseases	
Black stain root disease	Douglas-fir
<i>Armillaria</i> root disease	Douglas-fir, oaks, tanoak
Madrone canker	Madrone
Sudden Oak Death	Primary hosts on SDSF are California bay laurel and tanoak
Insects	
Flatheaded fir borer	Douglas-fir
Douglas-fir beetle	Douglas-fir
Redwood bark beetle	Coast redwood
Western oak bark beetle	Oaks, tanoak
California oakmoth	Oaks, tanoak
Mammals	
Tree squirrels	Coast redwood

There may be other pests of local tree species that are seldom detected or reported, or are of minor significance. State Forest staff will continue to monitor the Forest for early signs of forest pests or conditions that may lead to infestation.

Other efforts to reduce pest damage or predisposition will include:

- * The minimization of injuries to residual trees during forest management activities.
- * Reuse of old tractor roads or recreational trails where available to reduce soil compaction.
- * Retention of a diverse species composition in or adjacent to stands following forest management activities and within or nearby future regeneration units.
- * Avoidance of non-native tree species which may be predisposed to pests with few local pest predators and parasites.
- * Use of CAL FIRE or other forest pest management specialists to train SDSF staff in

forest pest recognition and management.

SUDDEN OAK DEATH (SOD):

California Code of Regulations--The Oak Mortality Regulation, as Applied to State Lands.

SDSF is within the declared SOD Zone of Infestation (ZOI) established by the California State Board of Forestry and Fire Protection and is within the “Regulated Area” for SOD as designated by the California Department of Food and Agriculture. The ZOI and Regulated Area are identical and cover all portions of the fourteen infested counties identified in the California Department of Food and Agriculture (CDFA) Section 3700: Oak Mortality Disease Control.

Federal regulations from the USDA-APHIS, and state regulations from CDFA address SOD concerns. CDFA regulations limiting the movement of host materials apply to forest management activities on SDSF including timber harvest, timber stand improvement activities, and harvest of minor forest products. Under the state regulation, host material cannot be transported from the Regulated Area unless accompanied by a compliance agreement. The Forest Practice Rules (FPRs) do not specifically address SOD. However, wherever a Zone of Infestation applies, the FPRs [14 CCR 917.9(a)] require that mitigations be included in Timber Harvesting Plans (THPs) to prevent the spread of the infestation. The following section includes a discussion of the applicable regulations, descriptions of host material, the “free-from” protocol, what constitutes a “compliance agreement,” and mitigation measures to prevent the spread of SOD.

A federal quarantine for *P. ramorum* was issued as an interim rule by USDA –APHIS, with the most recent rule dated February, 2007. For more information on the Federal rule see <http://nature.berkeley.edu/comtf/pdf/APHIS-2005-0102-0001.pdf>.

Host Material, The Oak Mortality Regulation

This is a California state law (Section 3700 in Title 3 of the Code of Regulations). The law defines the regulated articles (plants or plant parts) and commodities (unprocessed wood, wood products, and any other product, article, or conveyance presenting a risk of spreading the pathogen). On CDFA’s regulatory web page, with a link to APHIS’ updated February 2010 host list, 46 proven hosts and 81 associated plants (nursery stock) are now regulated as either entire plants or specific plant parts thereof. To review this list and keep abreast of updates, consult CDFA’s web site at <http://pi.CDFA.ca.gov/pqm/manual/htm/455.htm>.

Based upon currently available information, CDFA, the County Agricultural Commissioners, USDA Forest Service, and CAL FIRE have ranked the regulated articles according to their potential for transport of *P. ramorum* and believe that the highest risk for infected vegetation transport is from nursery stock of host species, followed by green waste, then firewood and logs. Lumber and manufactured wood products are not considered a risk for transport of *P. ramorum*. Soil is not included in California's regulations, but is in the federal regulations. Infested watercourses are not enforced in either. The enforcement rules are strictest for the highest risk regulated articles. Green waste is considered higher risk than firewood because the pathogen sporulates readily on leaves from tanoak, rhododendron, bay, and some other hosts. However, sporulation is rare on wood.

On SDSF, host material that is likely to be transported may consist of logs from host species produced as part of a harvest plan. In addition, there is the potential for minor forest products such as salvage sawlogs, firewood, and greenery to be moved from the regulated area.

Although SOD is present on the Forest, trained SDSF personnel will submit samples of symptomatic hosts and non-hosts to CDFA for confirmation of pathogen in new areas or on new hosts. As new hosts are confirmed by CDFA, the entire plant or specific portions thereof are amended to the rules as regulated articles. CDFA currently recognizes only RPFs, government agency personnel, and others that have been approved or attended specific CA Oak Mortality Task Force (COMTF)-certified training as "official samplers." The COMTF training session attendance roster may act as a list of official samplers.

Only an official sampler can complete a free-from survey as discussed below. The completed free-from survey, if part of an approved THP, will allow the plan to serve as a compliance agreement for a period of one year.

Regulated Area.

As of May 1, 2010, regulated counties are Alameda, Contra Costa, Humboldt, Lake, Marin, Mendocino, Monterey, Napa, San Mateo, San Francisco, Santa Clara, Santa Cruz, Solano, and Sonoma.

Distribution of SOD

Check the following website for known infested areas: <http://www.cnr.berkeley.edu/comtf/>. Information on local distribution may also be obtained from the County Agricultural Commissioner's office or CAL FIRE.

State Regulation.

SOD can spread via host material. Therefore, plants, plant parts, unprocessed wood and wood products, and other products of the above mentioned hosts cannot be moved from counties infested with SOD without authorization by the County Agricultural Commissioner or CAL FIRE's and USDA Forest Service's harvest document approval process. The term "harvest document" refers to any document filed with the California Department of Forestry and Fire Protection that authorizes the removal of forest products for commercial purposes. See <http://pi.CDFA.ca.gov/pqm/manual/htm/455.htm> for California Department of Food and Agriculture's (CDFA) regulations regarding commodities covered and restrictions of their movement.

Compliance Agreement.

A compliance agreement is required to move regulated articles of host material from the regulated area. A compliance agreement may be obtained through the County Agricultural Commissioner's office. An inspector will complete an inspection and make a determination that all the regulations are understood, that compliance with the regulations will be achieved and that

all provisions of the compliance agreement will be carried out. Once a landowner and USDA-APHIS sign a compliance agreement, the regulated host material may be moved interstate as stipulated in the compliance agreement.

A compliance agreement consisting of a free-from certification is only valid in California, as APHIS does not have allowance for free-from surveys in its regulations. A THP or other “harvest document” approved by CAL FIRE may also serve as a compliance agreement. An approved harvest plan without a free-from survey, but with approved SOD mitigation covering known hosts, will allow transport of host material from the regulated area. Approved plans with an incorporated free-from survey may act as a compliance agreement for a one-year period from the date of a documented negative result survey, or one year from the date of a negative reply from CAL FIRE to samples of symptomatic hosts.

Free-from Protocol

A *P. ramorum* "free-from" survey is a survey of land with host trees or shrubs that is done to determine if *P. ramorum* is present in the area in which a commercial operation will be conducted. All survey results are good for one year, unless symptomatic hosts are observed during that year. Transects would be run designed to cover representative areas of known SOD hosts. Transect width is variable to allow for adequate inspection. The official sampler must look for symptoms on all hosts along transects (also as approaching site, etc.). Samples will be sent to CDFA's Sacramento lab to confirm presence of *P. ramorum*. Since the survey is to determine presence or absence, if many symptomatic trees/shrubs are found, the survey may be stopped to wait for lab results. However, if *P. ramorum* is not found, the survey will need to be completed for the entire area.

Firewood sales on State Forest lands

Firewood sales for host material will not be permitted in areas with confirmed cases of sudden oak death (*P. ramorum*). Personnel setting up firewood sales must know how to recognize symptoms of *P. ramorum*. They should use the "free-from" protocol to determine if *P. ramorum* is present in the proposed firewood area. If *P. ramorum* is not detected in the "free-from" survey, the

operation is set up as usual with no further regulation for a period of one year. For firewood permits, a provision will be added to the permit explaining the current regulations, and that compliance is required.

On State Forest lands, where the public is required to have a permit to collect firewood, CAL FIRE shall use this public contact to educate the person(s) about sudden oak death by providing a Pest Alert or other information on sudden oak death along with the firewood permit.

For Timber Harvesting Plans

If host material is to be moved intrastate from the regulated area, one of three procedures shall be followed:

1. A survey to determine the area is "free-from" *P. ramorum* is completed by Registered Professional Foresters (RPFs) or other official sampler. Symptoms of *P. ramorum* will be surveyed for during the timber cruise or pre-sale layout done to develop the Timber Harvesting Plan. If no *P. ramorum* is detected, the survey design and results may be incorporated into the

- plan. If the one-year free-from period expires prior to or during timber operations, a new free-from survey must be conducted and amended to the plan. Or,
2. If a free-from survey results in positive SOD finds, mitigation measures shall be written into the plan. At a minimum, mitigations shall include current regulations. In support of 14 CCR 917.9, the RPF is encouraged to add additional measures or Best Management Practices to limit the spread or build-up. Best Management Practices for forestry may be found on COMTF's web site at: <http://nature.berkeley.edu/comtf/pdf/ForestryGuidelines.pdf>. Or,
 3. In lieu of a free-from survey, the RPF may assume that all known SOD hosts found on the plan area are infected, and appropriate mitigation measures shall be incorporated.

Forest Practice Rules Compliance

Pursuant to 14 CCR 917.9(a) for commercial harvest subject to the California Forest Practice Rules, and within the declared Zone of Infestation, the plans must identify feasible measures to mitigate adverse infestation or infection impacts during timber operations (PRC 4527).

Federal (APHIS) Enforcement Rules to Prevent Spread of SOD

Regulated articles may be transported interstate from the regulated area only if accompanied by a certificate/compliance agreement which verifies the regulated articles have been treated according to measures in the federal register.

Acceptable treatments are:

1. Wood products such as firewood, logs, or lumber must be free of bark.
2. Soil (nursery industry) that has not been in contact with SOD-infected hosts and is free of duff, or soil which has been heat-treated at 180F for 30 minutes.
3. Wreaths, garlands, and greenery dipped for one hour in water held at 160F. Bay leaves used in wreaths also may be vacuum/heat treated.
4. Green waste may move to energy generation facilities under limited permits issued by the local Agricultural Commissioner.

Hazard Reduction for SOD

Severely diseased or dying oaks and tanoaks with stem infections are often attacked by bark beetles (western oak bark beetle and ambrosia beetles) and infected with decay fungi such as *Hypoxylon* spp. Trees with these secondary insect attacks and decay fungi may be structurally weakened and should be removed if adjacent to high-use roads and trails.

In addition, trees killed by SOD may lead to increased surface and crown fuels, thus potentially influencing fire behavior. Rather than just a public safety issue, removal of dead and dying trees should also lessen the risk of surface and crown fires.

Further SOD Education

SDSF should make interested public parties aware that the COMTF keeps track of the latest information and links on sudden oak death at their web site, www.suddenoakdeath.org.

PITCH CANKER

SDSF staff will incorporate the most current best management practices as identified by the California Pitch Canker Task Force for controlling the distribution and spread of Pitch Canker. The following are management practices to be applied should pitch canker be identified on SDSF:

a. The timely removal and disposal of trees dying from pitch canker may help prevent the buildup of destructive beetles which can attack other trees, and can carry the pitch canker pathogen to uninfected trees. The disposal of pitch canker diseased material should be done on-site so as not to spread the disease to uninfested areas. Limbs and small pieces of wood may be left on-site or they may be chipped or burned. Logs cut from pitch canker diseased trees may be split for firewood for local use, but infected logs shall first have the stem cankered sections bucked from them. The remaining wood should be seasoned beneath a tightly sealed, clear plastic tarp to prevent the buildup of destructive insects. California Department of Forestry and Fire Protection Tree Note #3, Controlling Bark Beetles in Wood Residue and Firewood, provides specific guidelines for placing tarps over and around firewood. Do not stack pine firewood next to living pine trees or transport it to uninfested areas.

b. The distribution of the disease is discontinuous; thus, there are infested as well as uninfested areas within the ZOI. CAL FIRE pest management specialists continue to monitor for the disease. SDSF staff should report any symptomatic knobcone pines their CAL FIRE pest management specialist for determination of presence of pitch canker.

c. Directions for Registered Professional Foresters and Licensed Timber Operators:

- Know when you are working within an infested area.
- The California Department of Forestry and Fire Protection (CAL FIRE) has the authority to impose conditions on the commercial harvest of trees from timberland within the ZOI. Such actions are to be carried out on a case-by-case basis and depend upon the harvest operation's potential to contribute to disease spread. For all timber operations regulated by the Department, the Department must be informed if pitch canker is present within the operating area.
- Do not transport infected or contaminated material to areas that are free of the disease.
- When cutting or pruning a diseased tree, clean tools with a disinfectant before using them in uninfested areas. Lysol® is an effective sterilizer. Make sure that clients and co-workers are aware of these guidelines.

d. Directions for Firewood Cutters:

- SDSF personnel should be kept informed whether pitch canker is determined to be present on the Forest. Any suspect areas shall be avoided for firewood harvest until an official determination is made as to presence or not of the disease. At time of firewood permit issuance, SDSF personnel shall direct firewood cutters to disease free areas of the Forest. Information on pitch canker disease recognition and regulations shall be provided with the collection permit in the event pitch canker becomes present on the Forest.
- Tools and machinery that are used to cut trees with pitch canker disease **WILL BECOME CONTAMINATED** with the pitch canker fungus. There is little chance of spreading pitch canker if contaminated tools are only used on dead trees or on trees that are not pines. However, if contaminated tools or machinery will be used on living pines, the tools should be cleaned and sterilized before use on uninfected trees or in uninfested areas. Lysol® is a suitable sterilizer for this purpose. A logical alternative to repeated cleaning of equipment is to reserve one set of

equipment for use only in infested areas, and another set for use only in uninfested areas.

° Do not transport pine firewood out of infested counties (Santa Cruz County is an infested county). Sell pine firewood locally using local use guidelines.

e. Directions for Other Forest Product Harvesters:

° SDSF personnel, at time of collecting permit issuance shall direct collectors to disease-free areas of the Forest. Information on pitch canker disease recognition and regulations shall be provided with the collection permit.

° Collectors shall not be permitted to remove pine products from trees infested with pitch canker disease.

f. Directions for Reforestation in Areas Affected by Pitch Canker:

° Material for replanting should be as local in origin as possible to retain the genetic integrity of the local population.

° Option 1. The preferred strategy for reestablishing knobcone pine would be to allow natural regeneration to occur. Site improvement to encourage regeneration may be required where a dense overstory precludes the development of a seedling stand. Where natural regeneration does occur, it can be expected that pitch canker will eventually infect some or most of the young trees. However, the trees will vary in their susceptibility and some may sustain little or no damage. This is the least intrusive approach to reforestation, with the possibility that a level of pitch canker resistance will be attained that eventually provides the desired density of mature trees.

° Option 2. If option one is acceptable in principle but there is insufficient seed to produce a stand, locally collected seed could be introduced. By collecting seed from asymptomatic trees, there is more likelihood that some are resistant, and a certain percentage of the progeny will carry this trait as well. It is thought that a small percentage of resistant individuals may be sufficient to establish a stand. To diversify the seed source, it is recommended that seed be gathered from:

(1) multiple trees, at least 100 meters apart where possible, (2) trees of differing ages, and (3) different heights within the same tree. Trees that have been planted should be avoided, as they may be non-local in origin. In addition, trees with evidence of disease, especially western gall rust, should not be used as a seed source.

° Option 3. The least desirable measure for maintaining knobcone pine presence is to transplant known resistant seedlings from a reputable source. The potential loss of genetic integrity could result in a nonnative stand. However, there is currently no program identifying genetically resistant knobcone pines.

TREE SQUIRRELS

Tree squirrels may damage sapling redwoods by stripping the bark from upper stems. This causes partial or complete girdling, leading to either top kill or sapwood decay. As damage from squirrels is associated with dense second growth redwood stands, thinning is the only recommended management action. Where trees are more widely spaced, squirrel damage will generally be reduced.

PLANNED ACTIONS

1. Continually monitor the Forest for signs of pests and notify the CAL FIRE Forest Pest Management Specialist of any findings. Take action as needed to minimize or eliminate

any problems.

2. Thin dense stands of trees where necessary to maintain healthy growing conditions.
3. Provide applied forest pest research opportunities to interested agencies, institutions, or organizations.
4. Coordinate with Santa Cruz County in the detection of and protection against gypsy moths or other introduced pests.


WATERSHED ASSESSMENT

PHYSIOGRAPHY OF THE WATERSHED

SDSF is located almost entirely within the drainage of the East Branch of Soquel Creek, its landbase covering approximately 21 percent of the basin. For the most part, the Forest is underlain by fine grained sedimentary rocks that are highly weathered and easily eroded. The headwater channel of the East Branch follows the San Andreas Fault Rift Zone and is heavily disturbed from landslides triggered by earthquake activity. Hillslopes throughout the Forest are commonly steep and prone to large, deep-seated landslides. Even greater instability occurs in active inner gorge zones near stream channels. When compared to mass erosion events (e.g., large landslides), surface erosion of the loamy soils is of minor significance.

As stated in the Property Description chapter, the Soquel Creek watershed has a Mediterranean climate with cool, wet winters and warm, dry summers. Mean annual precipitation in the East Branch watershed varies from 30 to 46 inches and takes place primarily between November and April. Discharge records from the USGS stream gauging station located immediately upstream from the bridge in the town of Soquel show that the average discharge is 1.05 cubic feet per second per square mile. High intensity, long duration winter storms, however, produce extreme levels of runoff. Major floods occurred during December, 1955 and January, 1982, producing log jams and flooding in the town of Soquel (Lassette and Kondolf, 2003; see Figure 16). Singer and Swanson (1983) state that Soquel has a chance of being flooded once every seven to ten years.

Figure 16. Annual Peak Streamflow for Soquel Creek.



BENEFICIAL USES

The two main beneficial uses of the East Branch of Soquel Creek are cold water fisheries and water supplies for various purposes. The East Branch supports spawning and summer rearing

habitat for coho salmon and steelhead trout, but the number of returning fish has declined significantly in the past 40 years (refer to the Fisheries chapter). For example, in 1982 steelhead populations were only one-third to one-quarter of what they were thought to have been in the 1960s (Singer and Swanson, 1983). Coho salmon were thought to be extirpated from the Soquel Creek basin since 1968 (SCCRCD 2003, NMFS 2010), but 170 young-of-the-year fish were documented during surveys conducted in a stream reach below SDSF in August 2008. The major factors limiting anadromous fish populations are believed to be low summer flows, and limited rearing habitat due to siltation of pools and removal of woody debris. Large wood loading, floodplain connectivity, and estuary function were listed as poor for the Soquel Creek watershed in the draft Central California Coast (CCC) coho salmon recovery plan (NMFS 2010). The East Branch of Soquel Creek is considered to be the southern most extent of coho salmon in California.

The other primary beneficial use in the main stem of Soquel Creek is water supply. There are eight dams in the Soquel Creek watershed that impede or block anadromous salmonid migration (NMFS 2010). Additionally, numerous small diversions exist that provide water for domestic, agricultural, and industrial purposes. Two permitted domestic water supply systems exist in the main Soquel Creek watershed that utilize surface flow, but there are no permitted systems utilizing surface water in the East Branch basin. The East Branch Soquel Creek watershed assessment conducted in 1993 reported that there were six surface water diversions in the East Branch basin (Cafferata and Poole 1993). The largest diversion in the East Branch is operated by the Olive Springs Quarry.

Diversions and overdrafting of groundwater in low discharge summer months have dewatered portions of the East Branch [p.22] stream channel during drought years. This is likely to be a critical limiting factor for the fisheries resource in the lower part of the basin. After complaints by the city of Capitola in 1988, the State Water Resources Control Board concluded that a watermaster was needed to effectively regulate water use under low flow conditions, particularly during drought years, due to the complicated interrelationships of the water rights on Soquel Creek (Cafferata and Poole, 1993). To date, however, no watermaster has been appointed (Alley and Associates, 2004).

LAND USE HISTORY

The major types of land use in the East Branch watershed are timber management, recreation on public land, ranching, and residential development. The East Branch watershed's long history of timber harvesting began with clearcutting in the 1870's and continued into the 1940's. Timber was originally removed by oxen (1870-1895), then by steam donkey (1895-1930), and finally by crawler tractor. Following World War II, selective harvesting replaced clearcutting, continuing to the present day. Currently, nearly all of the old-growth timber has been harvested except for minor reserved groves and widely scattered suppressed trees. Second-growth harvesting has taken place on approximately 881 acres over the past 10 years. In recent years, cable yarding has been used along with tractor logging to selectively harvest timber in the basin. Approximately 20 percent of the basin is dominated by chaparral and has never been harvested.

Recreation in the form of hiking and mountain biking takes place primarily in The Forest of Nisene Marks State Park and SDSF (SDSF also allows horseback riding). Additionally, adjacent

property owners often allow friends and neighbors to utilize their lands as an access route to the park or SDSF. The Olive Springs Quarry, located near the East Branch channel, has produced sand and gravel from granitic rock exposed along the Zayante Fault for over 45 years. Residential development has occurred in the chaparral communities and has steadily expanded over the past 40 years. Over the whole Soquel Creek watershed, housing development is rated as moderate to high, with approximately 7,000 housing units present in the basin. Residential and commercial development is considered a very high threat to coho salmon (NMFS 2010).

CUMULATIVE WATERSHED EFFECTS ASSESSMENT

Cumulative watershed effects (CWE) can be defined as the physical and biological impacts that result from multiple land use disturbances over space and time. These impacts occur within and away from the locations of actual land use and are transmitted through the fluvial system. When considering CWE, it is appropriate to estimate how current and future projects, when combined with impacts from past activities, will influence beneficial uses present in the basin under review. Techniques to determine whether CWE are significantly adversely impacting beneficial uses have been developed, but are generally considered to be inadequate for varying reasons. Existing CWE assessment approaches mostly range from checklists or indices that are subjective but inexpensive and simple, to complex physically based models that have large data needs and are difficult to apply (Litschert 2009).

Watershed analyses and assessments are often used to evaluate cumulative watershed effects. For example, Berg et al. (1996) found that watershed analysis was the most suitable approach for assessing cumulative watershed effects in the Sierra Nevada. While a formal watershed analysis has yet to be completed for the Soquel Creek watershed, several watershed assessments and studies have been completed over the past 30 years that contribute a considerable amount of information regarding cumulative watershed effects. These studies include: Singer and Swanson (1983), Cafferata and Poole (1993), Santa Cruz County Resource Conservation District (2003), Alley and Associates (2003, 2004), Balance Hydrologics (2003), Greening Associates (2003), Pacific Watershed Associates (2003), and Lassetre and Kondolf (2003).

Cafferata and Poole's rapid landscape-level watershed assessment for the East Branch of Soquel Creek was conducted in 1992 and 1993. It is described below in considerable detail. Since specific harvest units had not been defined, the entire Forest served as the project area for this assessment. CWE assessment guidelines suggest that assessment areas must be large enough to detect past impacts and small enough to determine what the impacts of the proposed projects will be on the area. This assessment completed for the East Branch watershed evaluated the current condition of stream channels in the basin and determined how they would likely be altered with future timber operations.

The watershed assessment for SDSF was broken down into two main tasks: a stream channel inventory and a hillslope erosion/sedimentation evaluation. The stream channel inventory was completed in 1992 and illustrated the current condition of the channel network within the assessment area. Channel stability was rated with the U.S. Forest Service's Pfankuch Method, while channel condition was evaluated with CDF's Guidelines for Assessment of Cumulative Impacts (CDF, 1994). Due to limited personnel and time constraints, stream channels within SDSF were analyzed in greater detail than channels located in other areas of the basin (Poole, 1992).

The hillslope component of the CWE evaluation was completed in the summer of 1993. Quantitative estimates of erosion risk and erosion volumes associated with anticipated activities (e.g., timber harvesting and road building) were made based on the results of the Critical Sites Erosion Study (CSES; Lewis and Rice, 1989; Rice and Lewis, 1991). A selected number of random sample locations were installed on existing, planned, and abandoned roads, as well as past harvest areas. The resulting data was input into equations to estimate the risk of generating critical, or large, erosion sites (i.e., a large erosion event producing more than 100 cubic yards per acre). These equations are based on measurements made for slope, amount of hillslope or road curvature, soil color, and rock hardness.

On-site measurements were made to assess other sources of erosion not addressed by the CSES. Estimates were then made of the amount of sediment that could result from the approximated erosion. The significance of the estimated sedimentation was studied by comparing it with expectations of sediment yield for Soquel Creek. Since long-term sediment data from the Soquel Creek basin was not available, information from the San Lorenzo River, a similar, neighboring basin, was used. The methodology utilized for the hillslope erosion and sedimentation analysis was designed by Rice (1993) and was used in several parts of the state in the 1990's.

Locations in the East Branch assessment area that were found to have poor stability and channel conditions were Amaya Creek, Hinckley Creek, and the East Branch between Ashbury Gulch and the upper SDSF boundary. The reaches with the best ratings for stability (high-fair) and channel conditions were the upper part of the East Branch above SDSF and the main stem of Soquel Creek below the junction with the West Branch. The reaches of the East Branch between Ashbury Gulch and the junction with the West Branch were reported as having intermediate stability and channel conditions.

Summarizing the results for the 10 miles of the East Branch surveyed, fair channel stability was the most frequent finding (10% good, 67% fair, and 23% poor). This compares reasonably well to the fair/poor relative overall rating for sediment and log contribution assigned to the East Branch by Singer and Swanson (1983). The exception was the highly impacted reach along the San Andreas Rift Zone. Additionally, the major tributaries were severely degraded by heavy landsliding activity. Very large quantities of sediment are stored in these headwater tributaries, as well as in wide, vegetated terraces along the lower reaches of the East Branch. Observation of the East Branch channel revealed that riffle stretches generally had cobble and gravel sized particles protecting the stream bottom surface. Inspection of stable pools below Ashbury Gulch, however, showed that this habitat type contained significant percentages of fine sediment.

Large woody debris was not found to be blocking anadromous fish passage in the East Branch, but was limiting habitat use in the major tributaries. Large wood was generally lacking along most of the East Branch of Soquel Creek below Ashbury Falls, except where it had accumulated in a few fairly stable locations. Large woody debris is valuable because of the pool habitat for young salmonids that forms around it in gravel dominated stream systems.

Results of the hillslope evaluation indicate that the risk of generating critical erosion sites is not significantly different from the average for the rain-dominated portions of the northern Coast Ranges in California (Cafferata and Poole, 1993). Very large amounts of existing erosion were measured on a few of the plots, however, and existing erosion was found to be considerably higher than the amount estimated from the critical site equations. For example, one of the road

plots had a culvert that had carved an exceptionally large gully below its outlet, while another had large amounts of cut and fill slope erosion 50 feet from a Class II stream. Therefore, the potential for producing large erosion events is clearly evident in this terrain, even though the critical site equations did not generate exceptionally large estimates.

The terrain slope and distance to a Class II stream were used as an index of the proportion of erosion which would become sediment. The average sediment delivery was found to be much higher than is commonly assumed in forested watersheds. This is due to very high delivery from relatively few plots with extensive amounts of existing erosion. Long-term sediment data does not exist for Soquel Creek at the USGS gaging station; the nearest station with long-term discharge records and a reasonable sediment record is found on the San Lorenzo River at Big Trees State Park. A statistical analysis indicated that peak storm discharges on Soquel Creek were significantly related to those of the San Lorenzo River. The peak storm discharge/annual sediment yield relationship generated for the San Lorenzo River was then used to estimate the annual sediment yield at Soquel Creek.

Extreme natural variability in sediment yield for Soquel Creek was found for 1952 to 1990, with the range spanning several orders of magnitude. The estimate of mean annual sediment yield based on the stream discharge record for Soquel Creek ($\sim 1,500 \text{ yd}^3/\text{mi}^2/\text{yr}$) was somewhat higher than that calculated based on hillslope erosion plot measurements. Due to the assumptions involved in making these estimates, however, it is unlikely that these estimated sediment yields are significantly different. Both estimates of sediment discharge for Soquel Creek are high for forested areas and illustrate the erosive nature of the basin. Cafferata and Poole (1993) concluded that it was unlikely, however, that the limited timber harvesting and rebuilding of abandoned roads planned for SDSF would significantly elevate the erosion rate and have a significant adverse impact on the aquatic system. The anticipated sediment production from SDSF's timber production was found to be small compared to the range of variability in sediment flux observed in this basin.

More recent watershed assessment work in the Soquel Creek watershed can be compared to the results of the Cafferata and Poole (1993) rapid assessment. For example, Lassetre and Kondolf's (2003) much more detailed large wood study included the lower part of East Branch of Soquel Creek. They reported an average large wood loading for this reach of $0.005 \text{ m}^3/\text{m}^2$, which they stated lies at the low end of the range of observed values for North American streams (e.g., approximately 36 times lower than for old growth coast redwood forests in Humboldt County). Similarly, Alley and Associates (2003) reported that large wood was extremely scarce in Soquel Creek compared to other coastal streams recently surveyed.

Large wood loading in the East Branch of Soquel Creek is low due to extensive log removal efforts by Santa Cruz County from the 1950's to the 1990's. The Santa Cruz County RCD watershed assessment (SCCRCD 2003) concluded that the scarcity of large wood limits juvenile salmonid production throughout the Soquel Creek watershed. Similar to other studies, their assessment states that large wood is scarce in the middle part of the East Branch and recruitment is low for this channel reach.

Balance Hydrologics (2003) reported that Soquel Creek has experienced prolonged periods (up to 25 to 30 years) of disturbed watershed conditions over the past 150 years. These conditions, they found, were caused by both natural processes and anthropogenic activities with the later

likely resulting in an increase in the natural, base rate of sediment production in the watershed, although small relative to natural rates.

The Santa Cruz County Resource Conservation District (2003) watershed assessment project concluded that overall lateral channel stability of Soquel Creek was relatively high during the past 54 years with minor adjustment at several points of meander along the East Branch and mainstem. Major sediment production from natural sources was found to occur in both the East and West Branches and is most pronounced following earthquakes, large magnitude floods or forest fires, and is usually related to landsliding.

The SCCRCD (2003) assessment states that several factors appear to limit distribution and abundance of steelhead. These factors include passage impediments, poor spawning habitat quality (high proportion of fine sediment, number of constricting, steep riffles below spawning glides), low spring and summer baseflows, limited amount of escape cover (provided by instream wood, undercut banks, unembedded boulders, water depth itself), elevated water temperature, and limited water depth. Throughout the watershed, low baseflows and sedimentation limit the amount and quality of rearing habitat.

PWA (2003) inventoried the SDSF road network (18.2 miles) and found 82 sites with significant sediment delivery potential. Fifty seven of these sites were at stream crossings. Three crossing sites were identified as having a high treatment immediacy, with a potential sediment delivery of approximately 1,631 yds³.

CONCLUSIONS

In summary, it can be stated that the East Branch of Soquel Creek watershed is a highly sensitive basin. This is due to its inherent natural hillslope instability related to seismic forces, steep slopes, weak bedrock, and frequent high intensity rainfall. Impacts from past management activities, prior to modern California Forest Practice Rules, have contributed to degraded conditions in the basin, as has residential development.

The East Branch system is stressed by fine sediment moving downstream from tributaries, partially due to its close proximity to the San Andreas Fault Zone. This material is filling pools required for rearing habitat for listed anadromous salmonids. Lack of sufficient volumes of large wood and overdrafting of water in low flow summer months compounds this problem. As a result, the aquatic environment is sensitive to further degradation and timber operations must be carried out with extra caution.

Due to the sensitivity of the hillslopes and current channel conditions, special considerations are needed when planning forest management activities in the East Branch. Appropriate mitigations must continue to be utilized and remedial improvements implemented to repair existing problem areas, such as those suggested in the PWA (2003) report. If these considerations are employed, future timber sales can occur without significant adverse impacts to the beneficial uses of the basin.

MONITORING AND ENHANCEMENT

Ongoing management of SDSF will involve maintaining proper drainage along roads and trails by repairing culverts, water bars, and other drainage structures to reduce or prevent soil erosion and stream sedimentation. An active watershed remediation program will continue to be used that includes monitoring watershed conditions and implementing enhancement projects. Monitoring has included a forestwide inventory of stored sediments and active landslides, and mapping mass wasting hazards and surface erosion potential. Sediment risk-reduction projects have been rated for cost-effectiveness (PWA, 2003) and will be implemented in conjunction with timber operations according to their priority as available funding permits.

PUBLIC COMMENTS AND CONCERNS

Public comments regarding the East Branch watershed have focused on its distinction and sensitivity. Individuals are concerned that forest management activities will result in general watershed degradation, including sedimentation of streams and excessive hillslope erosion. This is particularly important to them in regard to steelhead numbers and habitat, and possible impacts upon their land during high intensity rainfall. As stated in this and other chapters, careful pre-project evaluation and measures to ensure the maintenance of watershed integrity are a priority for SDSF. The CWE assessment work discussed above and other studies assist the Forest staff to understand and manage for the sensitive nature of the basin.

MANAGEMENT GUIDELINES

1. Design streamside management zones that properly address the inherently unstable nature of the East Branch watershed in SDSF. This includes extending zone widths beyond the standards set by the California Forest Practice Rules, for Class I fish-bearing watercourses, as is appropriate in sensitive areas. These zones must provide for the long-term recruitment of large wood, protection of the stream channels and banks, stream shading, sediment filtration, nutrient input, microclimate control, floodplain function, and prevention of significant ground disturbance.
2. Place heavy emphasis on road design and maintenance, since roads generally produce the largest percentage of management-related sediment in forested watersheds. New seasonal and temporary roads should be outsloped to avoid concentrating water that could trigger landslides or transport sediment directly into flowing streams. Of foremost concern for new road construction is the avoidance of localized unstable areas. For mainline roads, road drainage structures and watercourse crossings must be adequately sized to ensure that the risk of failure is minimized. Roads no longer needed must be properly abandoned. Wet weather use for roads impacting flowing streams should not occur and an active winter maintenance program is needed to ensure that drainage structures are adequately functioning. (Refer to the Roads And Other Improvements chapter for more information.)
3. No tractor operations will be permitted on slopes which average more than 35 percent without site-specific evaluations of slope stability and erosion potential. This will depend

on the ability to mitigate such operations to levels of insignificance.

PLANNED ACTIONS

1. Continue to utilize a document that records data on all timber operations activities done in the Forest that could influence soil and water resources. This document, titled Timber Operations History, is part of ongoing watershed work. For example, information on road construction and harvesting dates as well as the harvesting systems and site preparation techniques utilized are recorded and mapped.
2. Implement the treatment priorities for high and moderate risk inventoried sediment sources in the Soquel Demonstration State Forest watershed assessment area developed by PWA (2003).
3. Evaluate the performance of each previously-implemented remediation project to determine the success in reducing the risk of large-scale sedimentation. Redesign and modify any project not meeting its intended objective. (Refer to Appendix C, Monitoring Plan.)

ARCHAEOLOGY

INTRODUCTION

An archaeological and historical field survey of SDSF was conducted during the summer of 1991. The chief investigator was Dr. Brian Dillon, a consulting archaeologist associated with the California State University at Northridge. In March of 1992, an archaeological and historical report was generated from the survey. In addition to study results, this report also includes information from an archival records research and an extensive oral history interview process. During the survey, Dr. Dillon and his crew discovered six archaeological sites within the State Forest boundaries: two prehistoric and three historic sites as well as one site with both prehistoric and historic features. For a detailed account of the archaeology and history of SDSF, refer to Archaeological and Historical Survey of Soquel Demonstration State Forest, Santa Cruz County, California (Dillon, 1992). Additional studies covering SDSF history and archaeology are listed in the References section of this management plan.

The significance of each site was determined by its archaeological and historical value, as outlined in state and federal guidelines. Significance, as defined by these guidelines, is based on uniqueness and preservation, with both considered in the determination of a site's value. Unique refers to how many other similar features exist (on other sites), while preservation refers to the condition of the features remaining on the site. A site is not considered significant if it, although unique, has been completely destroyed as there is nothing left to protect or study.

A confidentiality policy exists which limits public disclosure of sensitive archaeological and historical resources. Consequently, site locations in SDSF with moderate to high levels of significance will not be revealed to the general public. The confidentiality policy protects the resources from artifact collection, site excavation, and vandalism. The policy was approved by the State Historical Resources Commission under authority of Public Resources Code Section 5020.4 (c).

Since 1991, at least five archaeological surveys have been conducted on SDSF. Citations of these surveys—most of which were carried out for the preparation of timber harvesting plans (THPs) and included limited portions of the Forest—are included in the References section of this document. As a result of these surveys and the initial one conducted by Dr. Dillon, most of the Forest has been examined at least once for archaeological resources. Surveys conducted since 1991 have resulted in the identification and recording of additional archaeological features at two of the sites first identified during the 1991 survey, two additional prehistoric archaeological sites and nine historic-era sites. The following is a brief description of the sites recorded to date within SDSF.

PREHISTORIC SITES

The prehistoric era is believed to have begun on the central coast about 4,000 years ago. The most active times were during the Late Prehistoric Period, 1000 - 1600 AD, when hundreds of sites were established. The prehistoric sites found in SDSF are from this Late Prehistoric Period.

SITE DESCRIPTIONS

P-44-000328 (CA-SCR-296)

This site contains bedrock mortars and rock art. It may also contain subsurface deposits. The presence of petroglyphs - consisting of multiple cupules on a sandstone boulder - prompted Brian Dillon to declare this the first cupule site to be recorded on the Pacific Coast between Monterey and Marin Counties.

P-44-000329 (CA-SCR-297/H)

This site has both prehistoric and historic features. The prehistoric feature is a sandstone boulder with multiple mortars. The historic features is, according to Dillon, "...a split-redwood corral dating to the depression era of the 1930s."

CA-SCR-298

This site contains a sandstone boulder with a single mortar hole.

P-44-000544(CA-SCR-346)

This site of a concentration of mortars and cupules observed on the surfaces of several sandstone outcrops and a sandstone boulder located on a midslope bench.

P-44-000 ____ (CA-SCR-____) [Numbers not yet assigned]

This site consists of several mortars on the surface of a sandstone boulder. Although the site has been recorded, a Primary Number and Trinomial have not yet been assigned.

HISTORIC SITES

The Historic Period began along the central coast in 1769 and continues to the present day. The historic sites found in SDSF span from the Mexican Period, 1822 - 1848, to the Anglo-American Period, 1848 to present. The most recent site in the Forest, however, dates back to World War II.

SITE DESCRIPTIONS

CA-SCR-299H

This site contains the remains of a sawmill that was in use until the mid-1940's. The buildings surrounding the mill were bulldozed in the 1960's.

CA-SCR-300H

This site contains a sawpit that was likely used approximately 165 years ago to saw large logs into smaller sizes.

CA-SCR-301H

This site contains the remains of a resort and sulphur baths that were very popular from the late 1870's to the early 1920's.

P-44-000545 (CA-SCR-347H)

This linear site consists of a 5,650-foot long segment of old dirt road that corresponds in location to a road labeled "Old Spanish Ranch Road" on the "Official Map of the County of Santa Cruz" (Punnett Brothers, 1906). The road prism was probably built before 1898 and possibly as early as 1856. Historic features identified at various points along it include portions of split redwood post and barbed wire fence and remnants of an old fruit orchard located on an adjacent parcel of privately-owned land.

PUBLIC COMMENTS AND CONCERNS

Following completion of the archeological study in SDSF, members of the public expressed a desire to learn about Dr. Dillon's discoveries and their significance. Individuals were interested in research or other studies that might result from the findings as well as seeing the archeological sites. Eventually, all sites of moderate and high significance will be more thoroughly evaluated and, depending on the results, may be available for public viewing.

MANAGEMENT GUIDELINES

1. Heritage resources on SDSF will be protected, especially during planning and management activities including timber harvesting, recreation, and forestry education.
2. Sites with moderate to high significance value will be preserved and kept confidential, as per the State Historical Resources Commission and Public Resources Code. If, after thorough and careful study, it is determined that certain sites can endure limited public use, they may be made accessible to the community.
3. Educational opportunities, including the display of resilient historic features, will be incorporated into SDSF's Demonstration and Forestry Education programs.

PLANNED ACTIONS

1. Provide the opportunity for scientific study and research of all archaeological and historical resources. Researchers working with sites of moderate or high significance will be expected to uphold all confidentiality policies.
2. After extended study of all archeological sites, create interpretive material for those sites determined to be safe from public harm. Information may be made available through signs, brochures, and staff or docent-led tours of historical areas.

3. Research additional historic information including maps, photographs, written documents, and interviews.

ROADS AND OTHER IMPROVEMENTS

SDSF FACILITIES

The first development of roads within SDSF can be traced back to the mid-nineteenth century. According to local historian Ron Powell (Rancho de Palo de Yesca: the Soquel Augmentation Rancho, n.d.), the Santa Cruz Gap Turnpike (an arterial between Santa Cruz and the Santa Clara valley) was completed in May of 1856. This road included parts of what are now Hihn's Mill Road, Long Ridge Road, and Spanish Ranch Road. Further development of roads occurred when Frederick A. Hihn acquired the SDSF property in 1864.

In the 1870's, Hihn built Sulphur Springs Road to reach his Sulphur Springs Resort as well as a private road for logging and mill access. These roads, along with segments of the Santa Cruz Gap Turnpike, created what is now known as Hihn's Mill Road. This route extends from the Olive Springs Quarry to the parking area off Highland Way and originally included most of what is now Sulphur Springs Road. Additionally, logging by Jared and Seth Comstock in 1878 led to the construction of Comstock Mill Road and portions of Robinwood Lane.

The remainder of the roads in SDSF were constructed by the three timber companies that owned the property after Hihn. This later road construction started with the Monterey Bay Redwood Company in 1924, continued through the CHY Company ownership in the 1960's and 1970's, and ended with the Pelican Timber Company in the 1980's. Figure 15 shows the locations of all existing forest and access roads.

The only portion of SDSF that does not contain roads is the area south of Badger Springs to Santa Rosalia Ridge. This area is bordered by Sawpit Trail (see the Recreation chapter for information on trails) on the east, the East Branch of Soquel Creek on the northwest, and the Forest boundary on the south. Steam Donkeys were used to log this section prior to the introduction of crawler tractors, so no major roads or trails were built. Scars on the slopes above Badger Springs are still evident from this type of logging.

The only bridges associated with SDSF cross the East Branch at the Highland Way entrance and a short distance upstream from the entrance of Amaya Creek on Hihn's Mill Road. The Highland Way bridge is a narrow nine-foot wide railroad car bridge supported by earthen abutments. Future plans call for replacing this bridge with a wider structure. This bridge and the nearby parking area are located on the Burch property. The second bridge is a 90-foot-long rail flatcar bridge installed by CAL FIRE in ?.

LEGAL ACCESS

When the State acquired the major portion of the Pelican Timber Company's holdings along the East Branch in 1988 (refer to the Administration chapter for details), the property was inaccessible from county roads. The Forest is surrounded by private property on three sides and The Forest of Nisene Marks State Park (largely undeveloped) to the south. To ensure access, two deeded right-of-ways through private property were granted at the time of acquisition, providing for both administrative and public access.

The first right-of-way is through the Burch property to the east of SDSF. This route includes 0.7 miles of Hihn's Mill Road out to Highland Way, and provides the only public vehicular access to SDSF. (Vehicles are not allowed in the Forest but can drive to and park in a designated area just off of Highland Way.) The second right-of-way is through the CHY Company property to the southwest of SDSF and includes 1 mile of Hihn's Mill Road from the Forest boundary to the Olive Springs Quarry. Olive Springs Road, which connects with the major arterial of Soquel-San Jose Road, can then be accessed through the quarry for administrative purposes only.

DRIVABLE ROADS

All drivable roads in SDSF have been evaluated for safety and stability and are open to varying degrees of seasonal vehicular use. A number of old logging roads and constructed skid trails in the Forest, however, still need to be accurately mapped and have their suitability for reuse determined. Table 8 is a summary of the drivable roads in SDSF with information on the road name, length, and location.

PUBLIC ACCESS ROADS

County roads leading to SDSF access points include Olive Springs Road, Highland Way, and Eureka Canyon Road. Olive Springs Road provides reliable year-round administrative access through the Olive Springs Quarry. As mentioned above, this road connects with Hihn's Mill Road at the southwestern edge of the Forest. This route from the quarry has a rock surface and is generally open.

Highland Way, which is east of Soquel-San Jose Road, leads to the eastern portion of SDSF and, as previously mentioned, provides the only public vehicular access. Highland Way has been unreliable during recent winters, however, due to landslide closures. Eureka Canyon Road, which winds through the mountains from Corralitos, provides a longer alternate route to the Highland Way entrance. Eureka Canyon Road is typically used when closures block Highland Way but is also subject to slides. The Aptos Creek Fire Road, located within The Forest of Nisene Marks State Park, provides administrative access as well as a public entrance (by foot or bicycle) at the east end of Ridge Trail.

ROAD BUILDING AND MAINTENANCE

Road building and maintenance are critical elements of forest management. Forest roads are usually the largest potential source of management-related erosion and they provide the means to recreate, harvest timber, access emergency situations, maintain facilities, and patrol the Forest. Therefore, they should be in the best possible condition at all times. Culverts, water bars, rolling dips, and drainage ditches, structures which divert water away from or off of roads, need to be regularly inspected and repaired as necessary. Winter inspection is crucial as the majority of water and topsoil movement occurs during this rainy season. New roads or skid trails will not be constructed or old roads or skid trails rehabilitated without thorough evaluations of topography, intended use, soil stability, drainage capabilities, and construction costs.

Table 8. Drivable road segments of SDSF.

ROAD NAME	LENGTH	LOCATION
Hihn's Mill Road	7.0 miles	Highland Way to Olive Springs Road
Amaya Creek Road	2.0 miles	Hihn's Mill Road to Comstock Mill Road
Comstock Mill Road	0.7 miles	Robinwood Lane to private property
Sulphur Springs Road	1.5 miles	Hihn's Mill Road to Ridge Trail
Corral Trail	1.2 miles	Sulphur Springs Trail to the end of roaded portion of Corral Trail
Tractor Trail	1.3 miles	Hihn's Mill Road to end of roaded portion of Tractor Trail
Longridge Road	1.5 miles	Hihn's Mill Road to private property
Amaya Basin Road	0.7 miles	
Amaya Pond Road	0.5 miles	
Amaya Spillway Road	0.1 miles	
Lower Helipad Road	0.1 miles	
Shortridge Road	0.1 miles	

There are approximately 19 miles of existing or abandoned roads within SDSF (Refer to Figure 8 Timber Management chapter). About 22 miles of new roads are needed to complete the Forest road network; approximately seven miles of roads would be constructed, and one to two miles of old road would be abandoned during the next ten years. In addition, approximately 15 miles of road proposed for construction will be abandoned eventually.

California Forest Practice Rules specific to road building and maintenance will be applied during all forest management activities. This will be particularly emphasized during timber harvesting operations. These regulations, along with other site-specific strategies, will help minimize erosion and sediment delivery to watercourses, visual disturbance, and road construction. Generally, all construction or rehabilitation of roads will include outsloping of road surfaces, rolling dips, and a minimum of inside ditches to minimize hydrologic connectivity to stream channels. A road management plan will be developed which includes information from the 2003 PWA road inventory, which identified 82 sites with significant sediment delivery potential to watercourse channels. .

PUBLIC COMMENTS AND CONCERNS

Since the roads of SDSF will not be available for motorized use by the public, comments regarding them have been minimal. The issue of Forest access, however, is another matter.

While most of the access concerns described in the Administration chapter relate to the lack of public entrances, the inadequacy of existing public entrances is a problem as well. Individuals have expressed that the entrances are not only remote, but also difficult to maneuver in some cases. This is particularly true of equestrians who trailer their horses to the Highland Way parking area. As detailed in the Administration chapter and Management Guideline 1 below, SDSF is actively seeking new legal access points. New entrance areas will be designed so that multiple uses can be accommodated.

Another concern is that excessive road building will take place in the Forest. Though SDSF does not have a detailed road construction plan, California Forest Practice Rules place restrictions on the amount and types of roads that can be built in forested areas. It is the intent of the Forest to adhere to these regulations in addition to the examination of all projects and possible impacts on a site-specific basis.

MANAGEMENT GUIDELINES

1. Work with neighboring ownerships to maintain and improve access to the Forest. Continue working with the Olive Springs Quarry and the County of Santa Cruz to ensure that both ends of Hihn's Mill Road remain open.
2. All roads and other improvements shall be monitored and maintained in good condition. This will provide for safety and help prevent surface and mass erosion. Ben Lomond Conservation Camp and California Conservation Corps crews, CAL FIRE heavy fire equipment operators, and volunteers will help with road maintenance and repair, as appropriate.
3. New roads and other improvements will be developed as needed for access, resource protection, and forest management activities. High standards and compliance with California Forest Practice Rules for new logging road construction and existing road rehabilitation to minimize soil damage will be emphasized. New construction and maintenance methods will be researched and incorporated to demonstrate their effectiveness.

PLANNED ACTIONS

1. Develop a road management plan for SDSF. An inventory, assessment, and risk-rating of forest roads was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (Pacific Watershed Associates, 2003). An assessment of constructed skid trails will be completed when funds become available. (Refer to Appendix C, Monitoring Plan.)
2. Road conditions will be continually assessed to determine which segments are no longer suitable for use. Approximately one to two miles of old road beds will be abandoned over the next 10 years.
3. Investigate and, if possible, acquire additional administrative and public access to SDSF.

Alternative funding sources will be examined for these acquisitions as well as for construction and maintenance. (Refer to the Administration chapter for more information on access issues.)

4. Upgrade Hihn's Mill Road and the Highland Way entrance parking area for all-season use. This includes road rocking and replacing culverts at risk of failure with larger culverts. Revise the right-of-way agreement with Burch to improve forest access. Upgrade Longridge, Amaya Creek, and Sulphur Springs Roads for improved emergency access.
5. Widen and provide permanent abutments to the Highland Way entrance bridge in cooperation with Burch/Redwood Empire. Develop a permanent structure for crossing the East Branch via Hihn's Mill Road below the confluence with Amaya Creek.
6. Maintain trash racks above the inflow to culverts in the Forest. Inventory, assessment, risk-rating, and cost-effectiveness of repair of culverts was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (Pacific Watershed Associates, 2003).
7. Initiate a mile-post system for roads and trails to facilitate management and emergency response.
8. Review the location of all roads, landings, and skid trails on unstable areas by a Certified Engineering Geologist.

REFERENCES

LITERATURE

Anderson, K.R. 1995. A status review of the coho salmon(Oncorhynchus kisutch) in California south of San Francisco Bay. California Department of Fish and Game report to the Fish and Game Commission.82 p.

Alley, D.W. and Associates. 1992. Recommended monitoring and logging guidelines for the Soquel Demonstration State Forest related to aquatic resources and flood control. Unpublished report submitted to Thomas Sutfin, SDSF Forest Manager.24 p.

Alley, D.W. and Associates, 2003. 2003. Soquel Creek Salmonid Assessment and Enhancement Plan. In: Santa Cruz County Resource Conservation District, Soquel Creek Watershed Assessment and Enhancement Project Plan. Capitola, California. 182 p.

Alley, D.W. and Associates. 2004. 2004 Soquel Creek Lagoon management and enhancement plan update. Final Report prepared for the City of Capitola. Brookdale, CA. 172 p.

Arora, D. 1986. Mushrooms demystified. Ten Speed Press, Berkeley, California. 959 p.

Balance Hydrologics, Inc. 2003. Soquel Creek Watershed Assessment: Geomorphology and Baseflow Hydrology. In: Santa Cruz County Resource Conservation District. Soquel Creek Watershed Assessment and Enhancement Plan. Capitola, CA. 89 p. plus Tables and Figures.

Barnhart, R.A. 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest)--steelhead. U.S. Fish and Wildlife Service Biological Report 82(11.60). U.S. Army Corps of Engineers, TR EL-82-4. 21 p.

Bartlett, Richard D. and Patricia P. Bartlett, 2009. Guide and Reference to the Snakes of Western North America (North of Mexico) and Hawaii. University Press of Florida.

Bartlett, Richard D. and Patricia P. Bartlett, 2009. Guide and Reference to the Amphibians of Western North America (North of Mexico) and Hawaii. University Press of Florida.

Bartlett, Richard D. and Patricia P. Bartlett, 2009. Guide and Reference to the Turtles and Lizards of Western North America (North of Mexico) and Hawaii. University Press of Florida.

Benyus, J.M. 1989. Northwoods wildlife: a watcher's guide to habitats. NorthWord Press, Inc., Minocqua, Wisconsin. 365 p.

Berg, N.H., K.B. Roby, and B.J. McGurk. 1996. Cumulative watershed effects: applicability of available methodologies to the Sierra Nevada. In: Sierra Nevada Ecosystem Project; final report to Congress. Davis, CA: University of California–Davis, Centers for Water and Wildland Resources: 39–78. Vol. 3. Chapter 2.

Berlekamp, B. and T. Sutfin, 1995. Draft Fisheries management Plan. Soquel Demonstration State Forest, Soquel, California. 18 p.

Blazej, N, 1997. Education Study for Soquel Demonstration State Forest. Soquel Demonstration State Forest, Soquel, California. 92 p.

Breschini, Gary S., *The Indians of Monterey County*. Monterey County Archaeological Society, Carmel.

Broadbent, S. M., 1072. The Rumsen of Monterey: An ethnography from historical sources. *Contributions of the University of California Archaeological Research Facility* 14:45-94.

Cafferata, P. and C. Poole. 1993. Watershed assessment for the East Branch of Soquel Creek. Final report submitted to the California Department of Forestry and Fire Protection, Sacramento, California. 58 p.

California Department of Fish and Game, 2008. Complete List of Amphibian, Reptile, Bird and Mammal Species in California. Sacramento, CA. 25 p

California Department of Fish and Game, 2009. State and federally listed endangered and threatened animals of California. Department of Fish and Game, Biogeographic Data Branch, California Natural Diversity Database. Sacramento, California. 12 p

California Department of Forestry and Fire Protection (CDF). 1994. Guidelines for assessment of cumulative impacts. Sacramento, CA. 28 p.

California Department of Forestry and Fire Protection, 2002. Instream Temperature Monitoring 2002, East Branch of Soquel Creek, Fern Gulch Creek, and Amay Creek. Soquel Demonstration State Forest, Soquel, California. 10 p.

CDPR (California Department of Parks and Recreation) Office of Historic Preservation, 1989. Archaeological Resource Management Reports (ARMR): Recommended Contents and Format. *Preservation Planning Bulletin Number 4()*, Sacramento.

----, 1991. Guidelines For Archaeological Research Designs. *Preservation Planning Bulletin Number 5*, Sacramento.

----, 1995. *Instructions for Recording Historical Resources*. Sacramento.

CDPR (California Department of Parks and Recreation) State Historic Resources Commission, 1993 *Guidelines for the Curation of Archaeological Collections*. Sacramento.

California State Board of Equalization. Timber yield taxes for 1977-1993. Timber Tax Office, 134 D Street, Suite 303, Eureka, California.

CaliforniaHerps.com, A Guide to the Reptiles and Amphibians of California

Clark, Donald Thomas, 1986. *Santa Cruz County Place Names*. Santa Cruz County Historical Society, Santa Cruz.

Conway, S. 1976. *Logging practices: principles of timber harvesting systems*. Miller Reeman Publications, Inc., San Francisco, California.

Dillon, B. 1992. Archeological and historical survey of Soquel Demonstration Forest. CDF Archeological Reports, Number 6. 177 p.

Foster, Daniel G. and Mark V. Thornton, 2001. Management Plan for CDF's Historic Buildings and Archaeological Sites. Archaeological Report Number 22, California Department of Forestry and Fire Protection, Sacramento.

Good, T.P., R.S. Waples, and P. Adams (editors), 2005. Updated status of federally listed ESUs of West Coast salmon and steelhead. US Department of Commerce, NOAA Tech memo. NMFS-NWFSC-66, 598 p.

Greening Associates. 2003. Soquel Creek Riparian Vegetation Assessment. Santa Cruz County Conservation District. Soquel Creek Watershed Assessment and Enhancement Plan. Capitola, CA. 45 p. plus Appendices.

Greenlee, J.M. 1981. Santa Cruz County Fire History: (Maps--CA: J.M. Greenlee, 1981. 1:25,344). University of California, Santa Cruz Maps References 6, 1528, 55, 673.

Harrington, Mark R., 1942. Culture element distribution, XIX: Central California coast. *University of California Anthropological Records* 7(1):1-146.

Heizer, Robert F., Ed., 1973. The Costanoan Indians. *De Anza College, Local History Series* 18. Cupertino.

Hilton, S. and T.E. Lisle. Measuring the fraction of pool volume filled with fine sediment. Manuscript in preparation.

Holland, V.L., M. Hanson, D. Keil, and L. Kelly. 1992. Preliminary biological assessment of Soquel Demonstration Forest, Santa Cruz County, California. Contract File 4904, California Polytechnic State University Sponsored Programs. 48 p.

Hunter, M.L. 1990. *Wildlife, forests and forestry: principles of managing forests for biological diversity*. Regents/Prentice Hall, Englewood Cliffs, New Jersey.

Jones and Stokes Associates, Inc. 1995. Environmental impact report for the Soquel Demonstration State Forest General Forest Management Plan. Draft. July 1995. (JSA 94-146.) Sacramento, CA. Prepared for the California Department of Forestry and Fire Protection, Sacramento, CA.

----. 1996. Environmental impact report for the Soquel Demonstration State Forest General Forest Management Plan. Final. June 1996. (JSA 94-146.) Sacramento, CA. Prepared for the California Department of Forestry and Fire Protection, Sacramento, CA.

Jones, Terry L., Nathan E. Stevens, Deborah A. Jones, Richard T. Fitzgerald, and Mark G. Hylkema, 2007. The Central Coast: A Midlatitude Milieu. In *California Prehistory: Colonization, Culture, and Complexity*, edited by Terry L. Jones and Kathryn A. Klar, pp. 125-146. Altamira Press, New York.

Knopp, C. 1992. Monitoring plan for an assessment of indices for measuring the condition of cold water fish habitat. Unpublished report submitted to the North Coast Regional Water Quality Control Board as part of a contract with the California Department of Forestry and Fire Protection. 19 p.

Kroeber, A. L., 1925. Handbook of the Indians of California. *Smithsonian Institution, Bureau of American Ethnology Bulletin 78*. Reprint (1976); Dover Publications, New York.

Lassette, N.S. and Kondolf, G.M., 2003. Process Based Management of Large Woody Debris at the Basin Scale, Soquel Creek, California. Final report prepared for the California Department of Forestry and Fire Protection and Soquel Demonstration State Forest. University of California, Berkeley, Department of Landscape Architecture and Environmental Planning, Berkeley, California. 130 p.

Levy, Richard S., 1978. Costanoan. In California, edited by R. F. Heizer, pp. 485-495. *Handbook of North American Indians*, Vol. 8. W. C. Sturtevant, general editor. Smithsonian Institution, Washington, DC.

Lewis, J. and R.M. Rice. 1989. Critical sites erosion study, volume II: site conditions related to erosion on private timber lands in northern California. Final report submitted to the California Department of Forestry and Fire Protection. 95 p.

Linsley and Kraeger Associates, Ltd. 1992. Summary report on stream flow and rainfall data from gauges in the Soquel Creek Watershed, July 1990-March 1992. Los Gatos, California.

Lisle, T.E. and S. Hilton. 1992. The volume of fine sediment in pools: an index of sediment supply in gravel bed streams. *Water Resources Bulletin* 28(2): 371-383.

Litschert, S.E. 2009. Predicting cumulative watershed effects in small forested watersheds. Ph.D. Dissertation. Colorado State University. Fort Collins, CO. 208 p.

Manson, M.W. and J.A. Sowma-Bawcom. 1992. Geology, slope stability and earthquake damage in Soquel Demonstration State Forest. California Department of Conservation, Division of Mines and Geology. Open-file report 92-8. 65 p.

Margolin, Malcolm, 1978. *The Ohlone Way: Indian Life in the San Francisco-Monterey Bay area*. Heyday Books, Berkeley.

Mayer, K.E. and W.F. Laudenslayer, Jr., eds. 1988. A guide to wildlife habitats of California. Multiple-agency cooperative project for the California Department of Forestry and Fire

Protection. 166 p.

McNally, M. and R. Hester. 1993. Soquel Demonstration State forest recreation study final report. Center for Environmental Design Research, University of California, Berkeley. 97 p.

Milliken, Randall T., 1995. *A Time of Little Choice: The Disintegration of Tribal Culture in the San Francisco Bay Area: 1769-1810*. Ballena Press, Menlo Park, California.

Moratto, Michael J., 1984. *California Archaeology*. Academic Press, New York.

Morse, A. and Sutfin, T., 2003. 2002 Steelhead Trout Population Survey Report. Soquel Demonstration State Forest, Soquel, California. 21 p.

Morse, A. and Sutfin, T., 2004. 2003 Steelhead Trout Population Survey Report. Soquel Demonstration State Forest, Soquel, California. 21 p.

National Marine Fisheries Service (NMFS). 2010. Public draft recovery plan for central California coast coho salmon Evolutionary Significant Unit. National Marine Fisheries Service, Southwest Region. Santa Rosa, CA.

Orre, Edgar R., 2004. An Archaeological Survey Report for the Fern Gulch Timber Harvesting Plan, Santa Cruz County, California. Report on file (S-30692) at the Northwest Information Center of the California Historical Resources Information System, Sonoma State University, Rohnert Park.

Orre, Edgar R. and Scott Stephenson, 2007. Historical Resource Record for Corral Trail Fence Line [no site number assigned yet.].

On file at the Northwest Information Center of the California Historical Resources Information System, Sonoma State University, Rohnert Park.

Pacific Watershed Associates, 2003. Erosion Prevention Planning Project for County Roads and Roads in the Soquel Demonstration State Forest, in the Soquel Creek Watershed, Santa Cruz County, California. Summary Report prepared for the Santa Cruz County Resource Conservation District and the California Department of Fish and Game. Arcata, CA. 31 p. plus appendices

Payne, Stephen, 1978. *A Howling Wilderness: A History of the Summit Road Area of the Santa Cruz Mountains, 1850-1906*. Loma Prieta Publishing, Los Gatos, California.

Poole, C. 1992. Stream channel inventory for the East Branch of Soquel Creek. Unpublished report submitted as a Senior Thesis, Department of Environmental Studies, University of California, Santa Cruz. 50 p.

Powell, R., n.d. Rancho de Palo de Yesca: the Soquel Augmentation Rancho. Unpublished Soquel Demonstration State Forest Headquarters, Soquel, California. 168 p.

Punnett Brothers, 1906. *Official Map of the County of Santa Cruz*.

State Lands Commission *Grants of Land in California Made by Spanish or Mexican Authorities*. Sacramento.

Reavis, B., 1991. Status of California Steelhead stocks. *In* Pacific States Marine Fisheries Commission and Association of Northwest Steelheaders (organizers), International Symposium on Steelhead Trout Management. Portland, Oregon. P. 57-63

Rice, R.M. 1989. On-site effects: the necessary precursors of cumulative watershed effects. Unpublished paper presented at a Cumulative Impacts Training Session, Fort Bragg, California. 12 p.

Rice, R.M., and J. Lewis. 1991. [Estimating erosion risks associated with logging and forest roads in northwestern California.](#) *Water Resources Bulletin* 27(5): 809-818.

----. 1992. A guide to data collection and analysis in support of an appraisal of cumulative watershed effects. Unpublished report prepared for private timber companies in California. 29 p.

----. 1992. Erosion risk and cumulative watershed effects associated with forest management in the upper Butano watershed, with special consideration of effects on the Pescadero Marsh. Unpublished report prepared for Big Creek Lumber Company, Davenport, California. 40 p.

Rice, R.M. 1993. A guide to data collection and analysis in support of an appraisal of cumulative watershed effects. Unpublished report prepared for private timber companies in California. Arcata, CA. 29 p.

Santa Cruz County Resource Conservation District (SCCRCD). 2003. The Soquel Creek watershed assessment and enhancement project plan. Capitola, CA. 92 p.

Singer, S. and M.L. Swanson. 1983. Soquel Creek storm damage recovery plan: a reconnaissance level study with recommendations for watershed management. Unpublished report of USDA Soil Conservation Service, Aptos, California. 71 p.

State Emergency Assessment Team (SEAT). 2003 Summit Martin Fires State Emergency Assessment Team Report affecting watershed in Santa Cruz and Santa Clara Counties. Sacramento, CA 130 p.

State Lands Commission, 1984. *Grants of Land in California Made by Spanish or Mexican Authorities.* Sacramento.

Sturm, E., E. Gilbert-Horvath, J. Garza, and R. MacFarlane, 2009. Creation of a captive broodstock program for southern coho salmon (*Oncorhynchus kisutch*) ; results from the initial rearing and spawning of the first brood year. NOAA-TM-NMFS-SWFSC-436. March

Thomas, J.H. 1961. *Flora of the Santa Cruz Mountains of California.* Stanford University Press, Stanford, California. 433 p.

U.S. Department of Agriculture, Soil Conservation Service in cooperation with University of

California, Agricultural Experiment Station. 1980. Soil Survey of Santa Cruz County, California. 148 p.

USDI, National Park Service, 1991. How to Apply the National Register Criteria for Evaluation. *National Register Bulletin* 15. Washington, D.C.

University of California, Berkeley Center for Environmental Design Research, 1996. Draft Education and Recreation Master Plan for Soquel Demonstration State Forest. Berkeley, California. 75 p.

Walsh, J. McL., 1976. John Peabody Harrington: The man and his California Indian fieldnotes. *Ballena Press Anthropological Papers* 6. Ramona, California.

PERSONAL COMMUNICATION

Anderson, T. and J. Brown. July, 1994. Habitat Surveyors, California Polytechnic State University.

Hope, D. January, 1993. Resource planner, County of Santa Cruz.

Mason, P. June, 1992. Local resident.

McCaslin, M. August, 1993. Representative of the Monterey Bay Salmon and Trout Project.

McCrary, B. May, 2003. Owner, Big Creek Lumber Company.

Smith, J. February, 1993. Biologist, San Jose State University.

Wilson, N. July, 1993. Representative of the Fungus Federation.

APPENDIX A: TEXT OF ASSEMBLY BILL 1965

ASSEMBLY BILL NO. 1965

CHAPTER 1338

An act to add Article 4 (commencing with Section 4660) to Chapter 9 of Division 4 of the Public Resources Code, relating to state forests.

(Approved by Governor September 29, 1987.
Filed with Secretary of State September 29, 1987.)

LEGISLATIVE COUNSEL'S DIGEST

AB 1965, Farr. Soquel Demonstration State Forest.

Under existing law, the Department of Forestry and Fire Protection is authorized to engage in the management, protection, and restoration of state forests in accordance with plans approved by the State Board of Forestry and Fire Protection.

This bill would declare it the policy of the state to establish the Soquel Demonstration State Forest in Santa Cruz County. The department would be responsible for the development and establishment of the state forest and for maintenance and operations. The bill would authorize the department to permit a limited amount of commercial timber operations within the forest in order to provide funds for its maintenance and operation. The bill would require the department, in conjunction with a specified advisory committee, to adopt a general plan for the park by January 1, 1989, and would require the general plan to be approved by the advisory committee prior to adoption by the department.

The department's duties and authority under the bill would only arise if the state acquires the property comprising the demonstration forest.

The people of the State of California do enact as follows:

SECTION 1. Article 4 (commencing with Section 4660) is added to Chapter 9 of Division 4 of the Public Resources Code, to read:

Article 4. Soquel Demonstration State Forest

4660. It is hereby declared to be the policy of the state to establish and preserve an intensively managed, multifaceted research forest which is representative of forest activities as a living forest in Santa Cruz County within northern California's coastal redwood belt. The coast redwoods, as the dominant tree species in this area, are a valuable natural resource and are unique in North America for their beauty, abundance, diversity, and public accessibility, and their extreme beauty and economic value requires special measures for their protection for the use, enjoyment, and education of the public.

It is the intent of the Legislature, in establishing the Soquel Demonstration State Forest, to provide an environment that will do all of the following:

- (a) Provide watershed protection for local communities and base-line monitoring and studies of the hazards, risks, and benefits of forest operations and watersheds to urban areas.
- (b) Provide public education and examples illustrating compatible rural land uses,

including sustained yield timber production, as well as the historic development of timbering and forestry machinery, within the context of local community protection and nearby pressures.

(c) Provide a resource for the public, environmental groups, elected officials, environmental planners, the educational community, and the media as an open environment for the inspection and study of environmental education, forestry practices, and effects thereof.

(d) Protect old growth redwood trees.

4661. The department may permit a limited amount of commercial timber operations on the property within the Soquel Demonstration State Forest in order to provide funds for the maintenance and operation of the state forest and to allow fulfillment of the objectives of Section 4660. Income from the state forest property shall sustain all costs of operation and provide income for research and educational purposes.

4662. The department is responsible for the development and establishing of the Soquel Demonstration State Forest and for ongoing maintenance and operations. The director shall appoint an advisory committee to assist the department in planning future management of the forest. The advisory committee shall include representatives of the Santa Cruz County Board of Supervisors, the Department of Parks and Recreation, the Board of Forestry and Fire Protection, the Forest of Nisene Marks Advisory Committee, and the Department of Fish and Game.

4663. The department, in coordination with the advisory committee, shall adopt by January 1, 1989, a general plan for the state forest which reflects the long-range development and management plans to provide for the optimum use and enjoyment of the living forest, as provided in Section 4660, as well as the protection of its quality and the watershed within the Santa Cruz area. The general plan shall be approved by the advisory committee prior to adoption by the department.

4664. The duties and authority of the department pursuant to this article shall only arise if the state acquires the property comprising the Soquel Demonstration State Forest.

APPENDIX B: SPECIES LISTS FOR SDSF

FAUNA OF SDSF

From Preliminary Biological Assessment of Soquel Demonstration State Forest, Santa Cruz County, California (Holland et al., 1992) and David Suddjian, Biological Consultant.

SCIENTIFIC NAME

COMMON NAME

FISH

<i>Lampetra tridentata</i>	Pacific lamprey
<i>Oncorhynchus mykiss iridius</i>	steelhead trout
<i>Oncorhynchus mykiss</i>	rainbow trout
<i>Gasterosteus aculeatus</i>	northern threespine stickleback
<i>Cottus</i> sp.	sculpin

AMPHIBIANS

<i>Taricha torosa torosa</i>	California Newt, Coast Range Newt
<i>Taricha granulosa granulosa</i>	Rough-skinned Newt
<i>Dicamptodon ensatus</i>	Pacific Giant Salamander,
	California Giant Salamander
<i>Batrachoseps attenuatus</i>	California Slender Salamander
<i>Ensatina eschscholtzii xanthoptica</i>	Yellow-eyed Ensatina
<i>Aneides flavipunctatus niger</i>	Santa Cruz Black Salamander
<i>Aneides lugubris</i>	Arboreal Salamander
<i>Bufo boreas halophilus</i>	California Toad, Western Toad
<i>Pseudacris sierra</i>	Sierran Tree Frog, Pacific
	Chorus Frog
<i>Rana boylei</i>	Foothill Yellow-legged Frog
<i>Rana draytonii</i>	California Red-legged Frog

REPTILES

<i>Actinomyx marmorata pallida</i>	Western Pond Turtle, Southern Pacific Pond
	Turtle
<i>Sceloporus occidentalis boucurtii</i>	Western Fence Lizard, Coast Range Fence
	Lizard
<i>Plestiodon skiltonianus skiltonianus</i>	Skilton's Skink, Western Skink
<i>Elgaria coerulea</i>	
<i>coerulea</i>	Northern Alligator Lizard, San
	Francisco Alligator Lizard
<i>Elgaria multicaranata multicaranata</i>	
	Southern Alligator Lizard, California
	Alligator Lizard

<i>Charina bottae</i>	Northern Rubber Boa
<i>Diadophis punctatus amabilis</i>	Pacific Ring-necked Snake
<i>Contia tenuis</i>	Sharp-tailed Snake
<i>Pituophis catenifer catenifer</i>	Pacific Gopher Snake
<i>Lampropeltis getula californiae</i>	California Kingsnake
<i>Lampropeltis zonata multifasciata</i>	California Mountain Kingsnake, Coast Mountain Kingsnake
<i>Thamnophis sirtalis infernalis</i>	California Red-sided Garter Snake, Red-spotted Garter Snake
<i>Thamnophis elegans terrestris</i>	Western Terrestrial Garter Snake, Coast Garter Snake
<i>Thamnophis atratus atratus</i>	Santa Cruz Garter Snake
<i>Coluber constrictus flaviventris</i>	Western Yellow-bellied Racer, Western Racer
<i>Crotalus oreganus oreganus</i>	Northern Pacific Rattlesnake, Western Rattlesnake

BIRDS

<i>Ardea herodias</i>	great blue heron
<i>Butoridesvirescens</i>	green heron
<i>Aix sponsa</i>	wood duck
<i>Mergus merganser</i>	common merganser
<i>Anas platyrhynchos</i>	mallard
<i>Cathartes aura</i>	turkey vulture
<i>Pandion haliaetus</i>	osprey
<i>Accipiter striatus</i>	sharp-shinned hawk
<i>Accipiter cooperii</i>	Cooper's hawk
<i>Buteo jamaicensis</i>	red-tailed hawk
<i>Buteo lineatus</i>	red-shouldered hawk
<i>Aquila chrysaetos</i>	golden eagle
<i>Falco columbarius</i>	merlin
<i>Callipepla californica</i>	California quail
<i>Actitis macularia</i>	spotted sandpiper
<i>Columba fasciata</i>	band-tailed pigeon
<i>Zenaida macroura</i>	mourning dove
<i>Megascops kennicottii</i>	western screech owl
<i>Bubo virginianus</i>	great horned owl
<i>Glaucidium gnoma</i>	northern pygmy-owl
<i>Asio otus</i>	long-eared owl
<i>Aegolius acadicus</i>	northern saw-whet owl
<i>Calypte anna</i>	Anna's hummingbird
<i>Selasphorus rufus</i>	rufous hummingbird
<i>Selasphorus sasin</i>	Allen's hummingbird
<i>Ceryle alcyon</i>	belted kingfisher
<i>Melanerpes formicivorus</i>	acorn woodpecker
<i>Sphyrapicus nuchalis</i>	red-naped sapsucker
<i>Picoides pubescens</i>	downy woodpecker

<i>Picoides villosus</i>	hairy woodpecker
<i>Colaptes auratus</i>	northern flicker
<i>Sphyrapicus ruber</i>	red-breasted sapsucker
<i>Empidonax difficilis</i>	Pacific-slope flycatcher
<i>Sayornis nigricans</i>	black phoebe
<i>Myiarchus cinerascens</i>	ash-throated flycatcher
<i>Contopus borealis</i>	olive-sided flycatcher
<i>Contopus sordidulus</i>	western wood-peewee
<i>Cypseloides niger</i>	black swift
<i>Chaetura vauxi</i>	Vaux's swift
<i>Tachycineta thalassina</i>	violet-green swallow
<i>Stelgidopteryx serripennis</i>	northern rough-winged swallow
<i>Hirundo pyrrhonota</i>	cliff swallow
<i>Corvus corax</i>	common raven
<i>Cyanocitta stelleri</i>	Steller's jay
<i>Aphelocoma coerulescens</i>	scrub jay
<i>Poceile rufescens</i>	chestnut-backed chickadee
<i>Baeolofus inornatus</i>	oak titmouse
<i>Psaltriparus minimus</i>	bushtit
<i>Certhia americana</i>	brown creeper
<i>Sitta pygmaea</i>	pygmy nuthatch
<i>Troglodytes troglodytes</i>	winter wren
<i>Thryomanes bewickii</i>	Bewick's wren
<i>Chamaea fasciata</i>	wrentit
<i>Cinclus mexicanus</i>	American dipper
<i>Regulus calendula</i>	ruby-crowned kinglet
<i>Regulus satrapa</i>	golden-crowned kinglet
<i>Polioptila caerulea</i>	blue-gray gnatcatcher
<i>Toxostoma redivivum</i>	California thrasher
<i>Catharus ustulatus</i>	Swainson's thrush
<i>Catharus guttatus</i>	hermit thrush
<i>Turdus migratorius</i>	American robin
<i>Ixoreus naevius</i>	varied thrush
<i>Bombycilla cedrorum</i>	cedar waxwing
<i>Sturnus vulgaris</i>	European Starling
<i>Vireo solitarius</i>	solitary vireo
<i>Vireo huttoni</i>	Hutton's vireo
<i>Vireo gilvus</i>	warbling vireo
<i>Dendroica coronata</i>	yellow-rumped warbler
<i>Dendroica townsendi</i>	Townsend's warbler
<i>Dendroica occidentalis</i>	hermit warbler
<i>Dendroica nigrescens</i>	black-throated gray warbler
<i>Dendroica petechia</i>	yellow warbler
<i>Vermivora celata</i>	orange-crowned warbler
<i>Vermivora ruficapilla</i>	Nashville warbler
<i>Oporornis tolmiei</i>	MacGillivray's warbler
<i>Wilsonia pusilla</i>	Wilson's warbler
<i>Piranga ludoviciana</i>	western tanager

<i>Passerella iliaca</i>	fox sparrow
<i>Melospiza melodia</i>	song sparrow
<i>Melospiza lincolnii</i>	Lincoln's sparrow
<i>Zonotrichia atricapilla</i>	golden-crowned sparrow
<i>Zonotrichia leucophrys</i>	white-crowned sparrow
<i>Pipilo erythrophthalmus</i>	rufous-sided towhee
<i>Pipilo crissalis</i>	California towhee
<i>Junco hyemalis</i>	dark-eyed junco
<i>Pheucticus melanocephalus</i>	black-headed grosbeak
<i>Loxia curvirostra</i>	red crossbill
<i>Carpodacus purpureus</i>	purple finch
<i>Coccothraustes vespertinus</i>	evening grosbeak
<i>Carduelis tristis</i>	American goldfinch
<i>Carduelis psaltria</i>	lesser goldfinch
<i>Carduelis pinus</i>	pine siskin

MAMMALS

<i>Didelphis marsupialis</i>	opossum
<i>Sorex trowbridgei</i>	Trowbridge's shrew
<i>Neurotrichus gibbsii</i>	shrew-mole
<i>Scapanus latimanus</i>	broad-footed mole
<i>Myotis lucifugus</i>	little brown myotis
<i>Myotis evotis</i>	long-eared myotis
<i>Myotis volans</i>	long-legged myotis
<i>Myotis californicus</i>	California myotis
<i>Lasionycteris noctivagans</i>	silver-haired bat
<i>Lasiurus cinereus</i>	hoary bat
<i>Pipistrellus hesperus</i>	western pipistrelle
<i>Eptesicus fuscus</i>	big brown bat
<i>Sylvilagus bachmani</i>	brush rabbit
<i>Neotamias merriami</i>	Merriam's chipmunk
<i>Sciurus griseus</i>	western gray squirrel
<i>Thomomys bottae</i>	Botta's pocket gopher
<i>Chaetodipus californicus</i>	California pocket mouse
<i>Reithrodontomys megalotis</i>	western harvest mouse
<i>Peromyscus truei</i>	pinyon mouse
<i>Peromyscus californicus</i>	California mouse
<i>Peromyscus maniculatus</i>	deer mouse
<i>Neotoma fuscipes</i>	dusky-footed woodrat
<i>Urocyon cinereoargenteus</i>	gray fox
<i>Procyon lotor</i>	raccoon
<i>Mustela frenata</i>	long-tailed weasel
<i>Mephitis mephitis</i>	striped skunk
<i>Canis latrans</i>	coyote
<i>Puma concolor</i>	mountain lion
<i>Lynx rufus</i>	bobcat
<i>Sus scrofa</i>	feral pig

Odocoileus hemionus

black-tailed deer

FLORA OF SDSF

From Preliminary Biological Assessment of Soquel Demonstration State Forest, Santa Cruz County, California (Holland et al., 1992) and Randy Morgan. (Scientific names preceded by * are non-native.)

<i>Acer macrophyllum</i>	bigleaf maple
<i>Acer negundo</i>	California boxelder
<i>Achillea millefolium</i>	yarrow
<i>Adenocaulon bicolor</i>	trail plant
<i>Adenostema fasciculatum</i>	chamise
<i>Adiantum jordani</i>	California maidenhair
<i>Adiantum pedatum</i>	five-finger fern
<i>Aesculus californica</i>	California buckeye
<i>Ageratina adenophora</i>	sticky eupatorium
<i>Agoseris grandiflora</i>	large-flowered Agoseris
<i>Agrostis exarata</i> var. <i>pacifica</i>	Pacific bentgrass
* <i>Aira caryophyllea</i>	shiver grass
<i>Allium unifolium</i>	one-leaved onion
<i>Allophyllum divaricatum</i>	straggling gilia
<i>Alnus rhombifolia</i>	white alder
<i>Alnus rubra</i>	red alder
<i>Amsinckia intermedia</i>	common fiddleneck
* <i>Anagallis arvensis</i>	scarlet pimpernel
<i>Anaphalus margaritacea</i>	pearly everlasting
* <i>Anthemis cotula</i>	mayweed
<i>Anthriscus caucalis</i>	knotted hedge parsley
<i>Aquilegia formosa</i>	western columbine
<i>Aralia californica</i>	elk clover
<i>Arbutus menziesii</i>	madrone
<i>Arctostaphylos andersonii</i>	Santa Cruz mountain manzanita
<i>Arctostaphylos tomentosa</i> ssp. <i>crinita</i>	brittle-leaved manzanita
<i>Arctostaphylos</i> sp.	manzanita
<i>Artemisia douglasiana</i>	mugwort
<i>Asarum caudatum</i>	wild ginger
<i>Aster chilensis</i>	common aster
<i>Athyrium filix-femina</i>	lady fern
* <i>Avena barbata</i>	wild oats
<i>Baccharis douglasii</i>	marsh baccharis
<i>Baccharis pilularis</i> var. <i>consanguinea</i>	coyote brush
<i>Barbarea</i> sp.	wintercress
<i>Boisduvalia densiflora</i>	dense-flowered Boisduvalia
<i>Boykinia elata</i>	coast Boykinia
* <i>Brassica geniculata</i>	perennial mustard
* <i>Brassica nigra</i>	black mustard
* <i>Briza maxima</i>	rattlesnake grass
<i>Briza minor</i>	little rattlesnake grass
<i>Brodiaea elegans</i>	harvest brodiaea

<i>Brodiaea laxa</i>	Ithuriel's spear
<i>Bromus carinatus</i>	California brome
* <i>Bromus diandrus</i>	ripgut brome
* <i>Bromus hordeaceus</i>	soft chess
* <i>Bromus madritensis</i> ssp. <i>rubens</i>	red brome
<i>Bromus vulgaris</i>	common brome
<i>Calamagrostis rubescens</i>	pine grass
<i>Calochortus albus</i>	fairy lantern
<i>Calystegia occidentalis</i>	pine grass
<i>Campanula prenanthoides</i>	harebell
<i>Cardamine californica</i>	milkmaids
<i>Cardamine oligosperma</i>	popweed
* <i>Carduus pycnocephalus</i>	Italian thistle
* <i>Carduus tenuiflorus</i>	slender-flowered thistle
<i>Carex barbarae</i>	Santa Barbara sedge
<i>Carex bolanderi</i>	Bolander's sedge
<i>Carex globosa</i>	globose sedge
<i>Carex harfordii</i>	Hartford's sedge
<i>Carex tumulicola</i>	foothill sedge
<i>Ceanothus cuneatus</i>	buck brush
<i>Ceanothus papillosus</i>	warty-leaved ceanothus
<i>Ceanothus thyrsiflorus</i>	blue - blossom
* <i>Centaurea melitensis</i>	totalote
* <i>Centaurea solstitialis</i>	yellow star thistle
<i>Cerastium glomeratum</i>	mouse-eared chickweed
<i>Chlorogalum pomeridianum</i>	soap lily
<i>Cirsium brevistylum</i>	Indian thistle
* <i>Cirsium vulgare</i>	bull thistle
<i>Clarkia purpurea</i> ssp. <i>viminea</i>	large godetia
<i>Claytonia perfoliata</i>	miners' lettuce
<i>Collomia heterophylla</i>	Collomia
* <i>Conium maculatum</i>	poison hemlock
<i>Conyza canadensis</i>	Canadian horseweed
<i>Conyza sumatrensis</i>	Sumatran horseweed
<i>Convolvulus arvensis</i>	bindweed
* <i>Convolvulus occidentalis</i>	coast morning glory
<i>Cornus californica</i>	western red or creek dogwood
<i>Cornus sericia</i> ssp. <i>Sericia</i>	red osier dogwood
* <i>Cortaderia jubata</i>	pampas grass
<i>Corylus cornuta</i> var. <i>californica</i>	California hazel
* <i>Cynosurus echinatus</i>	dog's-tail grass
<i>Cynoglossum grande</i>	hounds tongue
<i>Cyperus eragrostis</i>	umbrella sedge
* <i>Dactylis glomerata</i>	orchard grass
<i>Danthonia californica</i>	California oat grass
<i>Dentaria californica</i>	milkmaids
<i>Deschampsia elongata</i>	hair grass
<i>Diplicis aurantiacus</i>	sticky monkeyflower

<i>Dichelostemma pulchellum</i>	blue dicks
<i>Disporum hookeri</i>	fairy bells
<i>Dryopteris arguta</i>	coastal wood fern
<i>Elymus glaucus</i>	blue wild rye
<i>Epilobium californicum</i>	California willow herb
<i>Epilobium ciliatum</i> var. <i>ciliatum</i>	willow herb
<i>Epilobium watsonii</i>	Watson's willow herb
<i>Epipactis helleborine</i>	eastern orchid
<i>Equisetum hyemale</i> ssp. <i>affine</i>	common scouring rush
<i>Equisetum telmateia</i>	horsetail
* <i>Erechtites glomerata</i>	fireweed
<i>Eriodictyon californicum</i>	California mountain balm
<i>Euphorbia peplus</i>	petty spurge
<i>Festuca californica</i>	California fescue
<i>Festuca occidentalis</i>	western fescue
<i>Festuca rubra</i>	red fescue
<i>Filago californica</i>	California Filago
<i>Fragaria californica</i>	California strawberry
* <i>Fragaria vesca</i>	woodland strawberry
<i>Fritillaria lanceolata</i>	checker lily
* <i>Galium aparine</i>	cleavers
<i>Galium californicum</i>	bedstraw
* <i>Galium murale</i>	tiny bedstraw
<i>Galium porrigens</i>	climbing bedstraw
<i>Galium triflorum</i>	sweet-scented bedstraw
* <i>Gastridium ventricosum</i>	nitgrass
* <i>Genista monspessulana</i>	French broom
<i>Geranium dissectum</i>	cut-leaved geranium
<i>Geranium molle</i>	dove's-foot geranium
<i>Gnaphalium californicum</i>	California everlasting
<i>Gnaphalium luteo-album</i>	weedy cudweed
<i>Gnaphalium purpureum</i>	purple cudweed
<i>Gnaphalium ramossissimum</i>	pink everlasting
<i>Gnaphalium</i> sp.	everlasting
<i>Gnaphalium stramineum</i>	cotton-batting plant
* <i>Hedera helix</i>	English ivy
<i>Helenium puberulum</i>	sneezeweed
<i>Heracleum maximum</i>	cow parsnip
<i>Heteromeles arbutifolia</i>	toyon
<i>Heuchera micrantha</i>	small-flowered heuchera
<i>Hieracium albiflorum</i>	hawk bit
<i>Hierochloa occidentalis</i>	vanilla grass
<i>Hirschfeldia incana</i>	mustard
* <i>Holcus lanatus</i>	velvet grass
<i>Holodiscus discolor</i>	cream bush
* <i>Hordeum leporinum</i>	foxtail barley
* <i>Hypericum calycinum</i>	St. John's wort
* <i>Hypochoeris radicata</i>	rough cat's-ears

<i>Iris fernaldii</i>	Santa Cruz Mountain iris
<i>Iris macrosiphon</i>	ground iris
<i>Juglans</i> sp.	walnut
<i>Juncus bufonius</i>	toad rush
<i>Juncus effusus</i> var. <i>brunneus</i>	bog rush
<i>Juncus effusus</i> var. <i>pacificus</i>	Pacific rush
<i>Juncus patens</i>	common rush
<i>Juncus xiphioides</i>	iris-leaved rush
<i>Lathyrus vestitus</i>	common Pacific pea
<i>Lathyrus vestitus</i> var. <i>vestitus</i>	wood-pea
* <i>Lepidium strictum</i>	wayside pepper grass
* <i>Linum bienne</i>	flax
<i>Lithocarpus densiflorus</i> var. <i>Densiflorus</i>	tanoak
* <i>Lolium multiflorum</i>	Italian wild rye
<i>Lonicera hispidula</i> var. <i>vacillans</i>	hairy honeysuckle
<i>Lotus eriophorus</i>	wooly trefoil
<i>Lotus micranthus</i>	trefoil
<i>Lotus purshianus</i>	Spanish clover
<i>Lotus scoparius</i>	deerweed
<i>Lupinus albifrons</i>	bush lupine
<i>Lupinus latifolius</i>	broad-leaved lupine
<i>Lupinus nanus</i>	sky lupine
<i>Luzula multiflora</i>	wood rush
<i>Madia elegans</i>	common Madia
<i>Madia gracilis</i>	slender Madia
<i>Madia madioides</i>	woodland Madia
* <i>Madia sativa</i>	common Madia
* <i>Marah fabaceus</i>	wild cucumber
* <i>Medicago polymorpha</i>	bur clover
<i>Melica imperfecta</i>	common melic grass
<i>Melica subulata</i>	Alaska onion grass
<i>Melica torreyana</i>	Torrey's melic grass
* <i>Melilotus alba</i>	white sweetclover
* <i>Melilotus indica</i>	yellow sweetclover
* <i>Melissa officinalis</i>	lemon balm
<i>Mimulus aurantiacus</i>	sticky monkeyflower
<i>Mimulus cardinalis</i>	scarlet monkeyflower
<i>Montia perfoliata</i>	miner's lettuce
* <i>Myosotis latifolia</i>	forget-me-not
<i>Myrica californica</i>	wax myrtle
<i>Navarretia squarrosa</i>	skunkweed
<i>Nemophila parviflora</i>	small-flowered Nemophila
<i>Nemophila pedunculata</i>	meadow nemophila
<i>Oemeria cerasiformis</i>	oso berry
<i>Osmorhiza chilensis</i>	sweet cicely
<i>Oxalis albicaris</i> ssp. <i>pilosa</i>	hairy wood sorrel
<i>Oxalis oregana</i>	redwood sorrel
<i>Oxalis pes-caprae</i>	sour grass

<i>Pentagramma traingularis</i>	gold-backed fern
<i>Petasites palmatus</i>	coltsfoot
<i>Petasites palmatus</i> var. <i>frigidus</i>	Western coltsfoot
<i>Phalaris californica</i>	California canary grass
<i>Pinus attenuata</i>	knobcone pine
<i>Piperia elongate</i>	rein orchid
<i>Piperia</i> sp.	royal rein orchid
<i>Pityrogramma triangularis</i>	goldenback fern
* <i>Plantago lanceolata</i>	English plantain
<i>Plantanus racemosa</i>	California sycamore
<i>Poa annua</i>	annual blue-grass
<i>Poa howellii</i>	Howell's blue-grass
<i>Polygala californica</i>	California milkwort
<i>Polygonum punctatum</i>	smartweed
<i>Polypodium californicum</i>	California polypody
* <i>Polypogon interruptus</i>	beard grass
* <i>Polypogon monspeliensis</i>	rabbitsfoot grass
<i>Polystichum dudleyi</i>	Dudley's shield fern
<i>Polystichum munitum</i>	western sword fern
<i>Populus trichocarpa</i>	black cottonwood
<i>Potentilla glandulosa</i>	sticky potentilla
<i>Prunus</i> sp.	wild plum
<i>Pseudotsuga menziesii</i>	Douglas-fir
<i>Psoralea macrostachya</i>	leather root
<i>Pteridium aquilinum</i> var. <i>pubescens</i>	bracken fern
<i>Quercus agrifolia</i>	coast live oak
<i>Quercus kelloggii</i>	black oak
<i>Quercus parvula</i> var. <i>shrevii</i>	Santa Cruz Mountain oak
<i>Quercus wislizenii</i>	interior live oak
<i>Rhamnus californica</i>	California coffeeberry
<i>Ribes menziesii</i>	canyon gooseberry
<i>Ribes menziesii</i> var. <i>senile</i>	Santa Cruz gooseberry
<i>Ribes sanguineum</i> var. <i>glutinosum</i>	pink-flowering current
* <i>Nasturtium officinale</i>	watercress
<i>Rorippa palustris</i>	yellow cress
<i>Rosa californica</i>	California wild rose
<i>Rosa gymnocarpa</i>	wood rose
<i>Rubus leucoderms</i>	black-capped raspberry
<i>Rubus parviflorus</i>	thimble berry
<i>Rubus ursinus</i>	California blackberry
* <i>Rumex acetosella</i>	sour dock or sheep sorrel
* <i>Rumex conglomeratus</i>	clustered dock
<i>Rumex salicifolius</i>	willow dock
<i>Rupertia physodes</i>	Rupert's scurf-pea
<i>Sagina apetala</i>	sticky pearlwort
<i>Salix lucida</i> ssp. <i>lasiandra</i>	yellow willow
<i>Salix lasiolepis</i>	arroyo willow
<i>Salix sitchensis</i>	velvet willow

<i>Sambucus mexicana</i>	blue elderberry
<i>Sanicula crassicaulis</i>	gambleweed
<i>Satureja douglasii</i>	yerba buena
<i>Scirpus microcarpus</i>	panicked bulrush
<i>Scoliopus bigelovii</i>	California fetid adder's tongue
<i>Scrophularia californica</i>	bee plant
<i>Sequoia sempervirens</i>	coast redwood
* <i>Silybum marianum</i>	milk thistle
<i>Sisyrinchium bellum</i>	blue-eyed grass
<i>Smilacina racemosa</i>	fat Solomon's seal
<i>Smilacina stellata</i>	slim Solomon's seal
<i>Solanum</i> sp.	solanum
<i>Solanum umbelliferum</i>	blue witch
<i>Soliva sessilis</i>	common soliva
* <i>Sonchus asper</i>	prickly sow thistle
* <i>Sonchus oleraceus</i>	smooth sow thistle
* <i>Spergularia</i> sp.	sand spurry
<i>Stachys bullata</i>	California hedge nettle
<i>Stachys chammisonis</i>	swamp hedge-nettle
<i>Stachys ajugoides</i> var. <i>rigida</i>	rigid hedge nettle
* <i>Stellaria media</i>	common chickweed
<i>Stipa pulchra</i>	purple needlegrass
<i>Symphoricarpos albus</i> var. <i>laevigatus</i>	snowberry
<i>Symphoricarpos mollis</i>	creeping snowberry
* <i>Torilis arvensis</i>	field hedge parsley
* <i>Torilis nodosa</i>	knotted hedge parsley
<i>Toxocodendron diversilobum</i>	poison oak
<i>Trientalis latifolia</i>	starflower
* <i>Trifolium angustifolium</i>	narrow-leaved clover
<i>Trifolium bididum</i> var. <i>decipiens</i>	pinole clover
<i>Trifolium dubium</i>	shamrock
<i>Trifolium gracilentum</i>	pin-point clover
* <i>Trifolium hirtum</i>	rose clover
<i>Trifolium microcephalum</i>	small-headed clover
<i>Trifolium obtusiflorum</i>	creek clover
<i>Trifolium wildenovii</i>	tomcat clover
<i>Trillium ovatum</i>	wake-robin
<i>Trisetum canescens</i>	tall trisetum
<i>Typha latifolia</i>	broad-leaved cattail
<i>Umbellularia californica</i>	California bay
<i>Urtica dioica</i> ssp. <i>holosericea</i>	hoary nettle
<i>Vaccinium ovatum</i>	huckleberry
<i>Vancouveria planipetala</i>	inside-out flower
<i>Verbena lasiostachys</i> var. <i>lasiostachys</i>	verbena
<i>Veronica americana</i>	American speedwell
<i>Veronica arvensis</i>	common speedwell
<i>Veronica peregrina</i>	purslane speedwell
<i>Veronica persica</i>	Persian speedwell

**Vicia sativa*

**Vinca major*

Viola ocellata

Viola pedunculata

Viola sempervirens

**Vulpia bromoides*

**Vulpia myuros*

Whipplea modesta

Woodwardia fimbriata

common vetch

greater periwinkle

two-eyed violet

Johnny jump-up

redwood violet

six-week fescue

rattail fescue

modesty

Western chain fern

FUNGI OF SDSF

<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>
<i>Agricus hondensis</i>	felt-ringed <i>Agricus</i>
<i>Amanita gemata</i>	gemmed <i>Amanita</i>
<i>Amanita pantherina</i>	panther <i>Amanita</i>
<i>Amanita phalloides</i>	death cap
<i>Amanita vaginata</i>	grisette
<i>Armillaria mellea</i>	honey mushroom
<i>Auriscalpium vulgare</i>	
<i>Boletus edulis</i>	king bolete
<i>Calodera viscosa</i>	yellow tuning fork
<i>Camarophyllus borealis</i>	snowy waxy cap
<i>Camarophyllus russocoriaceus</i>	cedar waxy cap
<i>Cantherellus californicus</i>	chanterelle
<i>Cantharellus cibarius</i>	
<i>Caulorhiza umbonata</i>	redwood rooter
<i>Clitocybe deceptiva</i>	anise mushroom
<i>Clitocybe nebularis</i>	cloudy <i>Clitocybe</i>
<i>Coprinus micaceus</i>	mica cap
<i>Cortinarius sp.</i>	<i>Cortinarius</i>
<i>Cortinarius varius</i>	<i>Cortinarius</i>
<i>Craterellus cornucopioides</i>	horn of plenty
<i>Crucibulum laeve</i>	white-egg bird's nest
<i>Entoloma sp.</i>	
<i>Fistulina hepatica</i>	beefsteak polypore
<i>Fomitopsis cajanderi</i>	
<i>Fomitopsis pinicola</i>	red-belted conk
<i>Galerina sp.</i>	
<i>Gomphidius subrosea</i>	rosy <i>Gomphidius</i>
<i>Gymnopilus sapineus</i>	common <i>Gymnopilus</i>
<i>Gyromitra infula</i>	hooded false morel
<i>Hebeloma crustuliniforme</i>	poison pie
<i>Helvella compressa</i>	
<i>Helvella lacunosa</i>	fluted black elfin saddle
<i>Helvella maculata</i>	
<i>Hemimycena sp.</i>	
<i>Heterotextus alpinus</i>	
<i>Hygrocybe acutoconica</i>	a cute conic waxy cap
<i>Hygrocybe coccinea</i>	righteous red waxy cap
<i>Hygrocybe conica</i>	witch's hat
<i>Hygrocybe flavescens</i>	golden waxy cap
<i>Hygrocybe punicea</i>	scarlet waxy cap
<i>Inocybe citrifolia</i>	
<i>Inocybe fastigiata</i>	corn silk <i>Inocybe</i>
<i>Lactarius argillaceifolius</i>	vulgar milk cap
<i>Lactarius chrysorheus</i>	yellow-staining milk cap
<i>Lactarius fragilis</i>	candy cap

<i>Lactarius rubrilacteus</i>	bleeding milk cap	
<i>Lenzites betulina</i>	gilled polypore	
<i>Leotia lubrica</i>		
<i>Leptonia parva</i>	blue-black Leptonia	
<i>Naematoloma fasciculare</i>	sulfur tuft	
<i>Paxillus involutus</i>	poison pax	
<i>Phylloporus rhodoxanthus</i>		gilled bolete
<i>Pleurocybella porrigens</i>	angel wings	
<i>Pleurotus ostreatus</i>	oyster mushroom	
<i>Pluteus cervinus</i>	deer mushroom	
<i>Psathyrella longipes</i>		
<i>Pseudohydnum gelatinosum</i>	jelly tooth	
<i>Ramaria sp.</i>	Ramaria	
<i>Ramaria gelatinosa</i>	jellied-base coral	
<i>Ramaria stricta</i>		straight-
<i>Russula brevipes</i>	branched coral	
<i>Russula cremoricolor</i>	short-stemmed Russula	
<i>Russula silvicola</i>	creamy Russula	
<i>Stereum hirsutum</i>	emetic Russula	
<i>Trametes versicolor</i>	hairy Stereum	
<i>Verpa conica</i>	turkey tail	
	thimble morel	

From personal communication with Nathan Wilson of the Fungus Federation of Santa Cruz

APPENDIX C: SOQUEL DEMONSTRATION STATE FOREST MONITORING PLAN

Identified monitoring projects will be implemented as staffing and budget limitations allow. To finance construction, these projects are generally incorporated as part of the timber harvest planning process.

WATERSHED MONITORING

TIMBER OPERATIONS

To avoid adverse impacts on water quality and fisheries resulting from the discharge of sediment to watercourses attributable to timber operations, Soquel Demonstration State Forest (SDSF) staff will monitor all timber operations (including all harvesting areas and new roads, skid trails, and landings) annually for 5-7 years following completion of the operations. Occurrences of substantial surface erosion (i.e., gullies) or mass wasting (i.e., landslides or slumps) resulting from the operations will be identified and described by a registered professional forester (RPF).

Each substantial gully or landslide will be evaluated to determine its cause and identify stabilization measures that would be most feasible, effective, and cost effective. Such measures will be implemented within 90 days from the date when the subject site is identified, unless due cause for delay is explained and a reasonable alternative schedule for implementation is proposed by the SDSF forest manager. If, based on the judgement of a certified engineering geologist (CEG), no stabilization measures are feasible or reasonable to apply to the subject site, feasible offsite watershed remediation measures will be implemented as recommended by the CEG in conjunction with the next timber operation conducted at SDSF.

The SDSF forest manager will describe appropriate mitigation measures to be incorporated into future timber operations and specified in future timber harvesting plans (THPs) to avoid a recurrence of the observed erosion or mass wasting events.

ROADS

An inventory that delineated, described, and risk-rated forest road at SDSF was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (Pacific Watershed Associates, 2003). This assessment included identification, , prioritization and design of feasible projects to remediate sediment-discharge risks associated with each high-and extreme-risk road segment and estimated the cost- effectiveness (in dollars per cubic yard of sediment) of each such project. Such projects include recontouring of roads and/or installation of improved drainage structures.

Soquel DSF staff conduct forensic monitoring of roads during and/or following most large storms. Any repairs needs are addressed as quickly as possible within the constraints of funding and legal and regulatory requirements.

CULVERTS

Soquel DSF staff evaluated the culverts along all Forest roads to determine where trash racks were necessary and appropriate. Steel T-posts were installed above each culvert where a need was determined. This work was completed prior to December 31, 1998. Each culvert is inspected annually and after large storm events, and cleaned as needed.

An inventory to delineate, describe, and risk-rate culverts at SDSF was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (Pacific Watershed Associates, 2003). This information was compiled into a database. The risk rating was based on standard engineering criteria for adequate culvert sizing, including effective culvert diameter, extent of tributary area, and intensity of a storm with a 100-year recurrence period.

The inventory included identification, prioritization and design of feasible projects to reduce the risk of failure of high-or extreme-risk culverts, and estimated the cost-effectiveness(in dollars per cubic yard of sediment) of each such project. Ppotential projects include replacement of existing culverts with more reliable drainage structures (e.g., rocked dips) or larger culverts.

Soquel DSF staff conduct forensic monitoring of drainage structures during and/or following most large storms. Any repairs needs are addressed as quickly as possible within the constraints of funding and legal and regulatory requirements.

ACTIVE LANDSLIDES AND SEDIMENT STORES

CAL FIRE, in conjunction with a CEG, will continue to inventory, delineate, describe and risk-rate active landslides and substantial unconsolidated concentrations of sediment and debris at SDSF. This information will be compiled into a database to be maintained on an ongoing basis with new landslides added as they occur. Unconsolidated concentrations of sediment and debris and active landslides identified by Manson and Sowma-Bawcom (1992) will be included in the database. The description and risk-rating of segments will focus on conditions affecting the likelihood that runoff or seismic activity will cause mass movement resulting in sediment discharge to a watercourse.

CAL FIRE , in conjunction with a CEG, will continue to identify and design feasible projects to reduce the risk of failure of high-or extreme-risk landslides and sediment-debris stores, and estimate the cost-effectiveness (in dollars per cubic yard of sediment) of each such project. Landslide risk-reduction projects could include installation of drainage structures on roads above landslides to reduce the discharge of concentrated runoff onto the landslide, or armoring or buttressing of landslide toes. Sediment-debris store risk-reduction projects could include recontouring or removal of material to reestablish original watercourses, bucking of logs in debris piles and using the logs to armor eroding bank faces, or revegetation.

AMAYA BASIN

The California Geological Survey compiled a map on landslides triggered by the 1989 Loma Prieta earthquake for Soquel Demonstration State Forest (Bedrossian, 1989). This map was focused on earthquake-triggered landslides and does not identify other unstable areas.

Additional reconnaissance-level slope stability mapping, which includes descriptions of the

geology and geomorphologic features related to slope stability was prepared by Manson and Sowma-Bawcom (1992), including all of the Amaya Creek and Fern Gulch watersheds. This mapping was based upon aerial photographic interpretations with little field work and is recognized as being useful for preliminary review of regional slope stability only.

In 2004, the California Geological Survey submitted an Engineering Geologic Report for the portion of Fern Gulch that was to be included in a proposed State Forest timber sale. This report includes a detailed landslide map designed to assist in timber sale development.

At the request of the Soquel Demonstration State Forest manager, the California Geological Survey began preliminary work for more detailed mapping of the Amaya Creek watershed in 2001, but this work was not funded and the mapping has yet to be completed.

REMEDICATION PROJECTS

CAL FIRE will prioritize all sediment-discharge projects identified and designed to remediate high- or extreme-risk conditions, as described above. Such projects will be implemented in conjunction with future timber operations according to their priority as available funding permits, to ensure, to the extent practicable, that no significant increase in sedimentation results from the timber operations proposed in each THP.

Soquel DSF staff monitor roads and drainage structures following most large storm events. When possible, inspections are conducted during storm events. Any repair needs are quickly addressed.

If CAL FIRE determines that any project is not meeting its intended objective, CAL FIRE will redesign and modify the project as needed.

AQUATIC RESOURCE MONITORING

FISHERIES MANAGEMENT PLAN

A Fisheries Management Plan for SDSF was completed in August, 1995. The aquatic-resource monitoring program includes the items described below.

FISH POPULATIONS

In cooperation with the California Department of Fish and Game (CDFG), fish population surveys were conducted at four separate sites on Soquel DSF from 1993 to 2001. From 2002 until the present, Soquel DSF has continued these annual surveys in cooperation with the National Oceanic and Atmospheric Administration's National Marine Fisheries Service (NOAA Fisheries). NOAA Fisheries added a fifth permanent station, increased the monitoring frequency, and incorporated additional research components.

HABITAT

One formal and comprehensive aquatic habitat survey was conducted on Soquel DSF in 1994.

Several additional surveys have been conducted by various fisheries biologists in conjunction with other research studies.

POOL SEDIMENTATION AND STREAM TEMPERATURE

The Aquatic Resources Monitoring Plan includes a pool sedimentation component to assess and monitor pool substrate embeddedness. At that time, the V* methodology was a popular assessment procedure to quantitatively measure sediment impairment. Fisheries biologists and hydrologists later advised SDSF staff that measuring sediment in a small number of pools, particularly using the V* method, would not be as useful as measuring other parameters, such as water temperature. To date, no formal survey has been conducted to assess pool sedimentation on watercourses within the Forest. The 1994 Fish Habitat Survey Report and other limited-scope investigations provide descriptions of the channel substrate composition.

On the advice of fisheries biologists, Soquel DSF staff initiated a long-term program to monitor stream temperature at various sites on the Forest. Temperature data has been recorded from 1997 through 2009.

AQUATIC INVERTEBRATES

A macroinvertebrate sampling study was performed by Soquel DSF staff in October, 1995. The California Department of Fish and Game analyzed the samples and produced a report of their findings in May, 1996. Since then, NOAA Fisheries has conducted additional assessments in conjunction with other research projects.

HABITAT ENHANCEMENT STRUCTURES

Fish habitat enhancement structures will be reviewed and approved by CDFG and an inter-agency team before installation.

To avoid adverse impacts of habitat-enhancement structures on fish habitat, CAL FIRE, in conjunction with CDFG, will evaluate structures annually for three years following installation to determine whether they are performing as intended and whether they are causing any unintended adverse impacts on fish habitat. If CAL FIRE and CDFG determine that any structure is functioning improperly or is adversely affecting aquatic habitat, CAL FIRE will redesign, repair, or remove the structure, as needed.

FISH POACHING AND HARASSMENT

To avoid adverse impacts of public use on fish, CAL FIRE and CDFG will conduct ongoing patrols in SDSF to enforce prohibitions on fishing and fish harassment. Incidents of fishing or harassment will be recorded and compiled. CAL FIRE and CDFG will evaluate such information annually in conjunction with fish population estimates to determine whether poaching or harassment have had a significant adverse effect on SDSF fisheries. If so, CAL FIRE will respond by closing streamside trails and intensifying law enforcement (e.g., increased patrols of streamside roads) as needed to ensure rapid fishery recovery and avoid additional

adverse fishery effects.

RIPARIAN RESOURCE MONITORING

To avoid adverse impacts of public use on riparian habitat, CAL FIRE will conduct ongoing patrols in SDSF to enforce prohibitions on vandalism and other damage to riparian habitat related to public use. Incidents of damage will be recorded and compiled. A qualified wildlife biologist will evaluate such information annually to determine whether the damage constitutes a significant adverse effect on wildlife. If so, CAL FIRE will implement additional restrictions on public use (e.g., prohibiting camping or weekday recreation use or cordoning off of sensitive areas) as needed to ensure rapid habitat recovery and avoid additional adverse wildlife effects.

WILDFIRE MONITORING

To minimize increases in wildfire risks resulting from increased public use at SDSF, CAL FIRE will record and compile descriptions of all wildfires occurring at SDSF including ambient weather and fire hazard conditions, ignition source, area and vegetation types burned, and estimated damage. Such information will be evaluated annually to determine whether wildfire incidents have increased substantially in frequency or intensity relative to 1990-1995 conditions. If so, CAL FIRE will implement appropriate measures (e.g., prohibiting all fires or weekday recreation use, or allocating additional fire-suppression resources for SDSF) as needed to reduce wildfire risks.

EMERGENCY SERVICES MONITORING

To minimize adverse effects on emergency response performance for residents of Santa Cruz County resulting from increased demands for such services at SDSF, CAL FIRE will record and compile all requests for emergency responses, including requests for police, fire medical, or search and rescue services. Descriptions of emergency responses will include response times. Such information will be evaluated annually to determine whether the demand for emergency services or average emergency response time has increased substantially relative to 1990-1995 conditions. If so, CAL FIRE will enhance its emergency response capability by upgrading roads; developing additional helispots along remote trails; and, either directly or through a management agreement with another qualified entity, providing additional human and equipment resources for emergency response at SDSF.

This data has not been compiled into a comprehensive report, but the data is examined by staff to evaluate any trends. All incident reports must be reviewed prior to public release to insure that legally confidential information (such as ongoing law enforcement actions, and the identities of juveniles or injured Forest visitors) is not disclosed. Soquel DSF staff do not have adequate resource to produce such a report.

NUISANCE MONITORING

To minimize nuisance impacts on SDSF neighbors, CAL FIRE will record and compile descriptions of all reported nuisances caused by SDSF users at SDSF or on adjacent ownerships including, but not limited to, trespass, vandalism, littering, and noise. This information has not been compiled into a comprehensive report, but the data is examined by staff to evaluate any trends.

Soquel DSF staff work closely with individuals and groups of volunteers to assist in this effort. The Stewards of Soquel Forest and a local National Mountain Bike Patrol group have both contributed information to the nuisance monitoring database. Soquel DSF staff have developed good relationships with neighbors to prevent incidents that constitute a nuisance.

PUBLIC-USE MONITORING

CAL FIRE will use camping records, surveys and other information to compile annual estimates of public use of SDSF in user days. Use will be estimated for specific activities including, but not limited to, mountain biking, camping, equestrian use, and educational activities.

CULTURAL RESOURCE MONITORING

CAL FIRE will monitor and periodically inspect heritage resources on SDSF to ensure that existing policies are affording effective protection. The identification and protection of cultural resources are important components of forestry in California today. Registered Professional Foresters are required to attend archeological training classes to acquire the ability to recognize cultural materials, and to be able to develop effective protection measures. In its role as a demonstration forest, SDSf can serve as a proving ground for the development and implementation of effective heritage resource management strategies.

CITATIONS

Manson, M.W., and J.A. Sowma-Bawcom. 1992. Geology, slope stability, and earthquake damage in Soquel Demonstration State Forest. California Department of Conservation, Division of Mines and Geology. Sacramento, CA.

Pacific Watershed Associates, 2003. Erosion Prevention Planning Project for County Roads and Roads in the Soquel Demonstration State Forest, in the Soquel Creek Watershed, Santa Cruz County, California. Summary Report prepared for the Santa Cruz County Resource Conservation District and the California Department of Fish and Game. Arcata, CA. 31 p. plus appendices

Figure 1. Locator map for SDSF.




Figure 2. SDSF and adjacent ownerships .




Figure 3. Locations of soil series in SDSF.




Figure 4. Vegetation Type Map for SDSF






Figure 5. Site Class Map for SDSF



SOQUEL DEMONSTRATION STATE FOREST



SOQUEL DEMONSTRATION STATE FOREST

LEGEND

- Forest Boundary
- Existing Roads
- - - Abandoned/Existing Roads
- - - Proposed Roads
- ▨ Old Growth Areas
- ▧ Non-Contiger Areas
- ▩ Cable Yarding Areas
- ▭ Tractor Yarding Areas




Figure 14. Recreational trails of SDSF and potential camp




Figure 15. Proposed fire defense improvements.




Figure 17. Drivable Roads of SDSF and Surrounding Area

